

**REPORT ON GEOMETRY AND REPRESENTATION THEORY OF TENSORS FOR
COMPUTER SCIENCE, STATISTICS AND OTHER AREAS, JULY 21 TO JULY 25,
2008 AT THE AMERICAN INSTITUTE OF MATHEMATICS, PALO ALTO,
CALIFORNIA, ORGANIZED BY JOSEPH LANDSBERG, LEK-HENG LIM, JASON
MORTON, AND JERZY WEYMAN**

LUKE OEDING

INTRODUCTION

This workshop was sponsored by AIM and the NSF and it brought in participants from the US, Canada and the European Union to Palo Alto, CA to work to translate questions from quantum computing, complexity theory, statistical learning theory, signal processing, and data analysis to problems in geometry and representation theory.

In all these areas, varieties in spaces of tensors that are invariant under linear changes of coordinates appear as central objects of study. Despite their different origins, there are striking similarities among the relevant varieties. We studied questions such as finding defining equations, hidden symmetries, and singularities.

The AIM workshop was preceded by a graduate student workshop at MSRI on “Geometry and Representation Theory of Tensors for Computer Science, Statistics, and other areas,” which was organized by Landsberg, Lim, and Morton. One goal of this workshop was to introduce the basic tools from geometry and representation theory which are used in studying problems that come up in these other areas. About half of the participants attended both workshops.

At AIM, the mornings were devoted to introductory talks given by L. Valiant, P. Comon, G. Gour, J.-Y. Cai, L. De Lathauwer, and J. Morton. In the afternoons the work focused on the following 4 areas, however many participants worked with more than one group.

- (1) Multi-linear techniques in data analysis and signal processing
- (2) Geometric approaches to $P?=NP$
- (3) Matchgates and holographic algorithms
- (4) Entanglement in quantum information theory

The participants split into these 4 groups, each of which contained a mix of practitioners and geometers. The groups then went to work on translating problems into a language which could be understood by both parties. After the translation work reached a stable point, the two goals were to determine what is already known in each area and to translate the open problems from the practitioners into well-posed problems in geometry. Finally, the participants made first steps toward answers - they determined simpler questions to start working on before the larger problems can be addressed; they made initial calculations to determine what should be true; and they made plans for collaboration focused on these problems.

ACKNOWLEDGMENTS

The author would like to thank A. Bernardi, A. Boralevi, P. Bürgisser, P. Comon, L. DeLathauwer, D. Gross, L. Gurvits, L.H. Kim, J.M. Landsberg, G. Ottaviani, and J. Weyman, for the many useful discussions both in person and via email during the preparation of this paper.

Part 1. Signal Processing

1. REPORT

The decomposition of tensors into a sum of rank one tensors has retained the attention of those in signal processing for quite a while. Collaboration with geometers continues to play an important role in determining what is already known and what tools and techniques may be immediately applicable to this and other problems in signal processing.

An important aspect of the workshop was developing a dictionary to translate notions in signal processing to notions in geometry. This dictionary opened the lines of communication between the engineers and the geometers. We will recap some of the relevant definitions here.

Let V_1, \dots, V_n be vector spaces (real or complex) and let $V_1 \otimes \dots \otimes V_n$ denote their tensor product. A *tensor* T is an element of $V_1 \otimes \dots \otimes V_n$. A *tensor decomposition* is the expression of a tensor as the linear combination of other tensors (presumably of lower rank). The *rank* of a tensor is the minimum r such that T may be expressed as the sum of r rank one tensors.

The rank of a tensor is unchanged by scalar multiplication, so it is natural to work in projective space. The following are the geometric objects that we will need.

The space of all rank one tensors is the Segre variety, defined by the embedding,

$$\begin{aligned} \text{Seg} : \mathbb{P}V_1 \times \dots \times \mathbb{P}V_n &\rightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n) \\ ([v_1], \dots, [v_n]) &\mapsto [v_1 \otimes \dots \otimes v_n]. \end{aligned}$$

The space of all rank one symmetric tensors is the Veronese variety, defined by the embedding,

$$\begin{aligned} v_d : \mathbb{P}V &\rightarrow \mathbb{P}(S^d V) \\ ([w]) &\mapsto [w^d]. \end{aligned}$$

The variety of partially symmetric rank one tensors is the Segre-Veronese variety defined by the embedding

$$\begin{aligned} \text{Seg}_{d_1, \dots, d_n} : \mathbb{P}V_1 \times \dots \times \mathbb{P}V_n &\rightarrow \mathbb{P}(S^{d_1} V_1 \otimes \dots \otimes S^{d_n} V_n) \\ ([w_1], \dots, [w_n]) &\mapsto [(w_1)^{d_1} \otimes \dots \otimes (w_n)^{d_n}]. \end{aligned}$$

Suppose $X \subset \mathbb{P}V$ is a variety. The r^{th} *secant variety* to X , denoted $\sigma_r(X)$, is the Zariski closure of all embedded secant \mathbb{P}^{r-1} 's to X , *i.e.*,

$$\sigma_r(X) = \overline{\bigcup_{x_1, \dots, x_r \in X} \mathbb{P}(\text{span}\{x_1, \dots, x_r\})} \subset \mathbb{P}V,$$

where the overline indicates Zariski closure. The *subspace varieties*, denoted $\text{Sub}_{r_1, \dots, r_k}(V_1 \otimes \dots \otimes V_k)$ are defined in [27] ch. 5. as

$$\text{Sub}_{r_1, \dots, r_k}(V_1 \otimes \dots \otimes V_k) = \{[T] \in \mathbb{P}(V_1 \otimes \dots \otimes V_k) \mid \forall i \exists A_i \subset V_i, \dim(A_i) = r_i, [T] \in \mathbb{P}(A_1 \otimes \dots \otimes A_k)\}.$$

(Note: these are related to “rank varieties” in [53] ch. 7.) We also consider the *symmetric subspace variety*,

$$\text{Sub}_r(S^n(V)) := \{[T] \in \mathbb{P}(S^n V) \mid \exists V' \subset V, \dim(V') = r, [T] \in \mathbb{P}(S^n V')\}.$$

In this setting, we list various notions of rank.

- * The *rank* of the point $[T] \in \mathbb{P}(V_1 \otimes \dots \otimes V_n)$ is the minimum r such that $[T]$ can be expressed as the sum of r points on the Segre variety.
- * More generally, for an algebraic variety $X \subset \mathbb{P}V$ the *X-rank* of a point $p \in \mathbb{P}V$, denoted $R_X(p)$, is defined by $R_X(p) := \min\{r \mid p \in \langle x_1, \dots, x_r \rangle, x_i \in X\}$, where $\langle \cdot \rangle$ denotes linear span.
- * The *X-border rank* is defined by $\underline{R}_X(p) := \min\{r \mid p \in \sigma_r(X)\}$. In particular, $\sigma_r(X)$ consists of all points with *X-border rank* no greater than r .
- * The *symmetric rank* of a tensor $[T] \in \mathbb{P}(S^n(V))$ is the *X-rank* when $X = v_n(\mathbb{P}V)$.
- * The *partially symmetric rank* of a tensor $[T] \in \mathbb{P}(V_1^{\otimes d_1} \otimes \dots \otimes V_k^{\otimes d_k})$ is the *X-rank* when $X = \text{Seg}_{d_1, \dots, d_k}(\mathbb{P}V_1 \times \dots \times \mathbb{P}V_k)$.

- * The *multilinear rank* of a tensor $[T] \in \mathbb{P}(V_1 \otimes \cdots \otimes V_n)$ is the minimum (r_1, \dots, r_k) such that $[T] \in \text{Sub}_{r_1, \dots, r_k}(V_1 \otimes \cdots \otimes V_n)$.
- * The *symmetric multilinear rank*, (deonted X_{sym} below) of a tensor $[T] \in \mathbb{P}(S^n(V))$ is the minimum r such that $[T] \in \text{Sub}_r(S^n V)$.
- * When no X is specified, the convention is that the *rank* (*border rank*) of a tensor $T \in V_1 \otimes \cdots \otimes V_n$ is the X -rank (respectively X -border rank) when $X = \text{Seg}(\mathbb{P}V_1 \times \cdots \times \mathbb{P}V_n)$.
- * For $X \subset \mathbb{P}V$, the *generic X -rank* is the X -rank of a generic tensor in $\mathbb{P}V$, *i.e.* it is the smallest r such that $\sigma_r(X) = \mathbb{P}V$.
- * A *typical X -rank* is an integer d such that the set $\{p \in \mathbb{P}(V_1 \otimes \cdots \otimes V_n) \mid R_X(p) = d\}$ has non-empty interior, *i.e.* it is an X -rank that occurs with nonzero probability. Note: over \mathbb{R} there can be more than one typical X -rank.

As far as applications are concerned, two important questions must be answered: (a) in what cases is a tensor decomposition unique, or at least when does there exist a finite number of such decompositions, and (b) how can one reasonably approximate a generic tensor by another of lower rank. A list of suggested problems from Comon (labeled P# below) were distributed before the workshop. During the workshop, the problems below were raised and discussed within the signal processing group.

- **The Alexander-Herschowitz theorem and generic ranks:** The Alexander-Herschowitz theorem describes the dimension of $\sigma_k(v_d(\mathbb{P}^N))$, which is the expected one, with a list of exceptions. This theorem is important because, in particular, it determines the generic symmetric rank of a tensor (see [3–5]). G. Ottaviani gave a nice exposition of the proof of Alexander-Hirschowitz, (see also [18, 41]).

The extension of the Alexander-Herschowitz theorem to not necessarily symmetric tensors should be the subject of future work (problem P5). It should be noted that progress in this direction has already been made, see [1]. Very recently, M. V. Catalisano, A. Geramita, and A. Gimigliano [9] have found that the secant varieties to $\text{Seg}(\mathbb{P}^1 \times \cdots \times \mathbb{P}^1)$ are all of the expected dimension with one exception.

- **Terracini’s lemma:** Terracini’s lemma (see [47]) is a tool which allows one to compute the generic rank numerically, when theoretical results do not allow one to obtain it. Terracini’s lemma is also often used as a theoretical tool in the Alexander-Herschowitz theorem, and the work of Chiantini-Ciliberto for example.

Terracini’s lemma also holds true for tensors with Hermitian symmetries, which is useful in Signal Processing when the data are complex (e.g. in Telecommunications) and when the tensor is estimated from the data. An example was presented by P. Comon regarding cumulant tensors. In the complex case, and for a given order, there are several cumulant tensors that can be defined. An example of an order 4 cumulant tensor is

$$T_{ijkl} = \text{Cum}\{x_i, x_j, x_k^*, x_l^*\},$$

where $(*)$ indicates complex conjugation. This one has plain symmetries $(ijkl \rightarrow jilk)$ and Hermitian symmetries $(ijkl \rightarrow klij)$ or $(ijkl \rightarrow lkji)$. Actually, these are the same symmetries as for the 4th order moment $E\{x_i, x_j, x_k^*, x_l^*\}$. See also [22].

- **Uniqueness of tensor decomposition:** Uniqueness of a given tensor decomposition is related to papers of Mella [36], and Chiantini-Ciliberto, [13, 14]. In the cases where the Alexander-Herschowitz theorem imply that there are at most a finite number of orbits, the work of Chiantini and Ciliberto gives a sufficient condition for uniqueness. Their condition is called weak defectiveness. This condition is straightforward, but to use it as a test is not a simple calculation. Mella gives an explicit list of some cases of uniqueness using different methods than Chiantini and Ciliberto. Certainly there is more work to do for the various types of tensor decompositions related to the other notions of rank mentioned above.
- **Real tensors (P10):** Tensor decomposition over \mathbb{R} is more complicated. In particular, there can be more than one typical rank, but it is not known if there can be more than two typical ranks. This

question has remained open after attempts by G. Ottaviani and P. Comon to find counter-examples. For the study of ranks in the real field, hyperdeterminants and other invariants should be useful.

- **Non-negative tensors (P11):** Tensors with real positive coefficients are useful in Spectral analysis and specifically to medical imaging. The approximation of a tensor by another of lower rank is well defined for tensors defined on the cone of non negative reals $\mathbb{R}_+^{\otimes n}$. A proof was sketched by L.-H. Lim.

The approximation of a tensor by another of lower rank is also well-defined when the vectors in at least one mode are constrained to be orthogonal. A proof was sketched by L. De Lathauwer.

A question that came up at the workshop is whether a specific metric is necessary. A metric is mandatory every time we have to deal with an approximation, which may happen for real positive, or complex, or real tensors. However, it was claimed that its exact form is not necessarily essential. The question of decomposing a non negative tensor into a sum of rank-1 terms can be posed without a metric. Whether a metric would help to solve the problem is another question. It may be that a topology is sufficient, as for the complex case.

- **Tensors with partial symmetries (P13):**

Consider the following types of tensors:

- The secant varieties to the Segre-Veronese
- 3-tensors enjoying the circular symmetries $T_{ijk} = T_{jki} = T_{kij}$.

These tensors are encountered in particular when using the joint characteristic function, estimated from data measurements. These are also related to the BIOME and FOABI algorithms.

(i) was discussed extensively during the workshop, whereas (ii) was only mentioned, but not focused on. It was pointed out that (ii) can be useful for the decomposition of the characteristic function, see [20], as well as other types of partial symmetries. The example of circular symmetry quoted above was mentioned by L. DeLathauwer. See also [21]

- **Usefulness of Kruskal's theorem for tensors:** Given an explicit decomposition of a tensor T , Kruskal's theorem provides a test as to whether that decomposition is unique. A translation into algebraic geometry terminology given by J.M. Landsberg allows one to shorten the proof. Possible generalizations to symmetric and partially symmetric tensors as well as extensions were discussed.
- **Comon's Conjecture (P15):** Do rank and symmetric rank always coincide? In other words, do we have equality in the relation

$$\sigma_k(\nu_d(\mathbb{P}V)) \subseteq \sigma_k(\text{Seg}(\mathbb{P}V \times \cdots \times \mathbb{P}V)) \cap \mathbb{P}(S^d V).$$

This has been proved for at least two cases: (i) $\text{rank}(T) \leq \text{f-rank}_{[1, \dots, k], [k+1, \dots, d]}(T)$ and $k < d/2$ (A proof of this fact, was derived and presented by D. Gross during the workshop. See Lemma 1.1 below for a definition of $\text{f-rank}_{[1, \dots, k], [k+1, \dots, d]}(T)$ as well as the proof of Gross.), and (ii) under Kruskal's condition $2 \text{rank}(T) \leq dn - d + 1$, where n denotes the dimension. See also [18].

For other cases, the problem remains open, despite attempts by A. Bernardi and L. Oeding to find counter examples.

- **Classification of varieties having rank equal to border rank:** We ask, for which varieties X is it true that the $R_X(p) = \underline{R}_X(p)$ for all $p \in \mathbb{P}(V)$?

Note that for any variety X , the tangential variety $\tau(X)$ is contained in the secant variety $\sigma(X)$. When X is the n -factor Segre variety, for instance, $\tau(X)$ contains points of rank n , however all of these points have border rank 2. This example shows that there can be an arbitrary discrepancy between rank and border rank.

- **Multilinear-ranks:** L. DeLathauwer introduced a generalization of block decomposition of matrices to block decomposition of tensors. The group pointed out links between this and subspace varieties. The geometric objects that encapsulate this idea are the secant varieties to the subspace varieties. We ask, what is the generic X -rank when X is the subspace variety? This question comes down to computing the dimensions of the secant varieties $\sigma_s(X)$.

We also want to consider symmetric multilinear rank. We ask, what is the generic X -rank when $X = \text{Sub}_r(S^k V)$? Already when $k = 3$ these are important questions with great interest to engineers in signal processing.

- **Strassen's Direct sum conjecture:** Suppose $X_1 = \text{Seg}(\mathbb{P}A_1 \times \cdots \times \mathbb{P}A_n)$ and $X_2 = \text{Seg}(\mathbb{P}B_1 \times \cdots \times \mathbb{P}V_n)$, and let $X = \text{Seg}(\mathbb{P}(A_1 \oplus B_1) \times \cdots \times \mathbb{P}(A_n \oplus B_n))$. Suppose $p \in (A_1 \oplus B_1) \otimes \cdots \otimes (A_n \oplus B_n)$. It is conjectured that $R_{X_1}(p) + R_{X_2}(p) = R_X(p)$.

Schönhage showed that this is false for border rank, so this seems to be a very subtle problem. P. Bürgisser gave a presentation of Schönhage's result.

- **Orbits (P6):** For which vector spaces with group actions do there exist finitely many orbits? In the cases that there are finitely many orbits, what are those orbits explicitly? J. Weyman pointed out that this problem has been solved. A paper of V. Kac has addressed the problem, [28], but there are mistakes, corrected in [29], pg 523, and in the multiplicity free case, [33].
- **Explicit decompositions** of tensors when the dimension is larger than two is an open problem (P7). Several suboptimal algorithms exist, for example [2, 21, 22, 31, 32], however they are limited in that the symmetries are not fully exploited and the rank must be smaller than some upper bound, hence, these algorithms cannot decompose generic symmetric tensors.

A suggestion was made by L. Gurvits to resort to doubly stochastic operators, but has not been pursued in depth.

- **Joint decomposition** of several tensors (P18). Considering two tensors (of possibly different orders) simultaneously may have two advantages: (i) There may be a possibility to restore uniqueness when the rank expected from the application exceeds the generic rank of the tensors (ii) There may be better robustness in the presence of noise. The problem was discussed (J.M. Landsberg, P. Comon, J. Weyman, G. Ottaviani), but postponed to further collaborations.
- **Sylvester theorem** was pointed out to be derived earlier by Gaspard de Prony (1755-1839), [23].

This is related to the signal processing problem of the estimation of frequencies of damped sinusoids.

It should be mentioned that more generally, all the questions raised for Segre or Veronese varieties (genericity, uniqueness, etc.) can be posed for subspace varieties.

For further reference for the ideas surrounding signal processing, tensor decompositions and questions of rank, we compiled the following list: [1, 10–12, 16, 17, 24, 30, 40, 46].

1.1. **From D. Gross.** Let $V = \mathbb{C}^n$ and $\text{Sym}^d(V)$ be the totally symmetric space in $V^{\otimes d}$. Let $T \in \text{Sym}^d(V)$. By a *decomposition* of T , we mean a set $D = \{a_i\}_i$, where

$$a_i = a_i^{(1)} \otimes \cdots \otimes a_i^{(d)}$$

is a rank-one tensor that

$$T = \sum_i a_i.$$

A decomposition D is *symmetric*, if all elements in D are symmetric. Let $I \subset \{1, \dots, d\}$. We write

$$\text{f-rank}_{I, I^C}(T)$$

for the rank of T viewed as a degree-2 tensor obtained by grouping the respective indices in I and I^C together (the 'f' stands for "forgetful" or "flattening"). Also, if D is a decomposition, write

$$D^{(I)} = \left\{ \bigotimes_{j \in I} a_i^{(j)} \right\}_i$$

for the projection of the decomposition onto the factors in I . We get

Lemma 1.1 (D. Gross). *Let $T \in S^d(V)$, $d > 2$. If, for any k ,*

$$\text{rank } T \leq \text{f-rank}_{[1, \dots, k], [k+1, \dots, d]}(T)$$

then any minimal decomposition of T is symmetric.

What is more, let $D = \{a_i\}_i$ be a decomposition of a symmetric tensor. Assume that for any $I \subset \{1, \dots, d\}$ where $|I| = d - 2$ the set $D^{(I)}$ is linearly independent. Then D is symmetric.

Proof. We prove the second assertion first. Let $I = \{1, \dots, d-2\}$. By assumption, $D^{(I)}$ is linearly independent, so there exists a dual basis $\{\alpha_i\}_i$ fulfilling $\alpha_i(a_j) = \delta_{i,j}$ for every $a_j \in D^{(I)}$. Hence, for every i ,

$$\alpha_i(T) = a_i^{(d-1)} \otimes a_i^{(d)}.$$

Since $\alpha_i(T) \in \text{Sym}^2(V)$, it follows that $a_i^{(d-1)} = a_i^{(d)}$. Now take $I = 1, \dots, d-3, d$. Arguing as before, we arrive at $a_i^{(d-2)} = a_i^{(d-1)}$. By induction $a_i^{(k)} = a_i^{(l)}$ for all k, l , proving the claim.

The first statement follows, because $\text{f-rank}_{I,I^c}(T) \leq \text{rank}(T)$ and equality trivially implies that $D^{(I)}$ is linearly independent for any minimal decomposition D . \square

Note that the first assertion is formulated in terms of geometric properties of T , whereas the second statement just limits our ability to write down non-symmetric decompositions of symmetric tensors.

2. OPEN PROBLEMS

- (1) Consider the case when $X = \text{Sub}_{r,\dots,r}(V \otimes \dots \otimes V)$, and $X_{\text{sym}} = \text{Sub}_r(S^n(V))$. Under what general conditions is it true that $X\text{-rank}(T) = X_{\text{sym}}\text{-rank}(T)$ for all tensors T in $S^n(V)$? We can ask the same question for border rank. Notice that when $r = 1$, $\text{Sub}_{1,\dots,1}(V \otimes \dots \otimes V) = \text{Seg}(\mathbb{P}V \times \dots \times \mathbb{P}V)$ and $\text{Sub}_1(S^n(V)) = v_n(V)$. Therefore, Comon's conjecture is a special case of this problem.

It was remarked that for $r = 1$ some cases are known. For instance if $\dim(V) = 2$ and arbitrary r , etc. See also, [19]. There are some very old results, for instance, Prony's method from 1795, which is equivalent to Sylvester's algorithm, [23].

- (2) Again consider the case when $X = \text{Sub}_{r,\dots,r}(V \otimes \dots \otimes V)$, and $X_{\text{sym}} = \text{Sub}_r(S^n(V))$. What is the generic X -rank and X_{sym} -rank for X and X_{sym} ? Note that r_1, r_2, r_3, r , etc, must have $r_i \leq r_j r_k$ for i, j, k distinct, otherwise it collapses.

Suggestion of Weyman: Suppose one would like to know about the existence a generic rank in this context. One can translate this to a question of the existence of an open orbit. There is a close relation to the Castling Transform. Start with $\mathbb{C}^p \otimes \mathbb{C}^q \otimes \mathbb{C}^r$ with $p \leq q \leq r$ and $r \leq pq$. Then transform the question to $\mathbb{C}^p \otimes \mathbb{C}^q \otimes \mathbb{C}^{pq-r}$. Then you can restrict to an open set and get a correspondence on orbits. Therefore the property of the existence of an open orbit occurs simultaneously for (p, q, r) and $(p, q, pq - r)$.

- (3) Let $X = v_k(\mathbb{P}V)$ and consider $Y_r = \{[T] \in S^k V \mid X\text{-rank}(T) \leq r\}$. Conjecture 6.10 [18] claims that $Y_r \neq \overline{Y}_r$ for $1 < r < R_S$, where R_S is the minimum r such that $Y_r = S^k V$. In particular, this says that the set of symmetric tensors of rank at most P is closed only for $P = 1$ and $P = R_S$.
- (4) What can we say about the uniqueness or finiteness of the corresponding decompositions $t = t_1 + t_2 + \dots + t_d$, where $t_i \in X$, and $X\text{-rank}(t) = d$. We can ask this question when X is any of the varieties we have mentioned. Particularly interesting are the cases when X is a subspace variety, or a symmetric subspace variety. Again, many cases are already known, for instance see [46].
- (5) Is the subspace variety version of Strassen's direct sum conjecture true? Let $t_1 \in A_1 \otimes B_1 \otimes C_1$, $t_2 \in A_2 \otimes B_2 \otimes C_2$, so that $t_1 \oplus t_2 \in (A_1 \oplus A_2) \otimes (B_1 \oplus B_2) \otimes (C_1 \oplus C_2) = A \otimes B \otimes C$. Let $X_i = \text{Sub}_{a_i, b_i, c_i}(A_i \otimes B_i \otimes C_i) \subset \mathbb{P}(A_i \otimes B_i \otimes C_i)$ and $X = \text{Sub}_{a,b,c}(A \otimes B \otimes C) \subset \mathbb{P}(A \otimes B \otimes C)$ be the respective subspace subvarieties. We ask if the following equality holds:

$$X\text{-rank}(t_1 \oplus t_2) = X_1\text{-rank}(t_1) + X_2\text{-rank}(t_2).$$

What, if any, relationship must there be between X, X_1 and X_2 ? What is the status of the conjecture for other varieties X, X_1, X_2 ?

- (6) The previous 4 questions are posed over the base field \mathbb{C} . Over \mathbb{R} , what are the typical X -rank's and X_{sym} -rank's in the semi-algebraic sense?
- (7) Tensor product of cones over orthants: Let $A_+ = \mathbb{R}_+^a, B_+ = \mathbb{R}_+^b, C_+ = \mathbb{R}_+^c$. Let $X_+ \subset A_+ \otimes_{\mathbb{R}_+} B_+ \otimes_{\mathbb{R}_+} C_+$ be a semi-algebraic variety. Study the X_+ -rank.

A tensor with all non-negative coefficients can be written as a non-negative linear combination of tensors with rank ≤ 1 . In this case we must have $X_+ = \text{Seg}_+ = (\text{Sub}_{1,1,1})_+$, and $X_+ = \text{Seg}(\mathbb{P}A \times \mathbb{P}B \times \mathbb{P}C) \cap V_+$. It was remarked that (of course) this rank can be larger than the X -rank.


Part 2. Mulmuley-Sohoni Approach to \overline{VP} vs. VP vs. VNP

3. REPORT

The Mulmuley and Sohoni approach to P vs. NP has as its primary goal to “reduce hard nonexistence problems in complexity theory to tractable existence problems in geometry and representation theory”, [39]. Before we get to this geometry and representation theory, we will briefly review some of the relevant notions.


During the workshop, L. Valiant described algebraic versions of the complexity classes P and NP (these are commonly referred to as VP and VNP), [51]. Computing the permanent is known to be an VNP -hard problem, however, by using techniques from linear algebra, computing the determinant can be computed in polynomial time. Valiant’s approach to P vs. NP suggests attempting to show that if one computes the permanent of an $m \times m$ as the determinant of an $n \times n$ matrix, that n must grow faster than a polynomial in m . (For reference, see [37, 38].)

P. Bürgisser, J.-Y. Cai pointed out that Toda defined a skew arithmetic circuit (see [48–50]). An arithmetic circuit can be described by a finite directed tree, where all of the edges (usually called wires) are directed towards the root of the tree, and each of the nodes (usually called gates) has precisely two incoming wires and one outgoing wire. At the top of the circuit, the input to each wire is either a constant or a variable. The wires take their inputs to gates, and each gate does a single elementary computations on those inputs. In an arithmetic circuit, there are only two types of gates allowed - multiplication and addition. So each gate takes precisely 2 polynomials as inputs and outputs either the sum or the product of the input polynomials. A skew arithmetic circuit carries the additional requirement that every multiplication gate have at least one of its inputs either a variable or a constant. Locally, a skew arithmetic circuit for multiplication looks like the following.



The inputs of this circuit are the variable x_i and the polynomial g and the output is their product, $x_i g$. The final output of an arithmetic circuit is a polynomial. Toda showed that the determinant of an $n \times n$ matrix can be computed by a skew arithmetic circuit whose size is bounded by a polynomial in n . A sequence of polynomials is in VP if it can be computed by a sequence of skew arithmetic circuits whose sizes and degrees are bounded by a polynomial in the number of inputs.

Toda also defined a notion of weakly skew. Malod and Portier, [34] recall that a weakly skew arithmetic circuit is an arithmetic circuit with the requirement that of the two sub-circuits above each multiplication gate, at least one must become disjoint from the rest of the circuit upon deletion of its wire to the multiplication gate. This may be depicted roughly as follows:



The boxes represent sub-circuits of the original circuit and $(*)$ is a multiplication gate. A sequence of polynomials is in VP_{WS} if it can be computed by a sequence of weakly skew arithmetic circuits whose sizes and degrees are bounded by a polynomial in the number of inputs. These are examples of Valiant’s complexity classes. It is not known whether $VP = VP_{WS}$ in general, however, Valiant comments that if you allow $n \log(n)$ - i.e. quasi-polynomial, then all of these classes collapse.

The sequence (det_n) is complete for the class VP_{WS} [34]. It is not known whether $VP = VP_{WS}$. Since the sequence $(perm_m)$ is VNP complete, Mulmuley and Sohoni are trying to prove that $\overline{VP}_{WS} \neq VNP$, where the class \overline{VP}_{WS} is defined analogously to \overline{VP} as in the paper of Bürgisser, [6].

Since we don’t want to be deceived by bad choices of coordinates, we work with the appropriate orbits of $perm$ and det . The Mulmuley-Sohoni approach to P vs NP is to determine if there exists a polynomial

$n(m)$ such that

$$[l^{n-m}perm_m] \in \overline{[GL(W).det_n]},$$

where $W = Mat_{n \times n} \mathbb{C}$. We also want to know what we can say about the inclusion

$$[End(W).det_n] \hookrightarrow \overline{[GL(W).det_n]}.$$

It should be noted that the description of the orbit closure of a given polynomial is in general very complicated.

The strategy suggested is to understand the homogeneous coordinate rings. We should try to understand the modules of functions $S \subset \mathbb{C}[\overline{GL(W).l^{n-m}perm_m}]$ which do not occur in $\mathbb{C}[\overline{GL(W).det_n}]$. Mumfley-Sohoni call such an S a *representation theoretic obstruction* for $l^{n-m}perm_m$ to lie in the variety $\overline{GL(W).det_n}$, [39].

Remark 3.1. Let V be a G -module for a reductive group G , and let $v \in \mathbb{P}V$. Then we have the following inclusions $G.v \subset \overline{G.v} \subset \mathbb{P}V$. The big problem is to understand the restriction $\mathbb{C}[\overline{G.v}] \hookrightarrow \mathbb{C}[G.v]$. The only general tool is a theorem of Kostant. It says that if the closure $\overline{G.v}$ is normal and $\overline{G.v} \setminus G.v$ has codimension 2, then the restriction map is an isomorphism. However in our examples these conditions are almost certainly not satisfied. Representation theory gives the following decomposition: Let G_v denote the stabilizer in G of v , then $\mathbb{C}[G.v] = \mathbb{C}[G/G_v] = \bigoplus_{\lambda \in Irrep(G)} V_{\lambda}^* \otimes V_{\lambda}^{G_v}$. This gives information on the coordinate ring of the orbit closure.

Step 1: The first example is if $v = [det_n]$, and $W = M_n \mathbb{C} = A \otimes B$, then

$$\mathbb{C}[GL(W).det_n] = \bigoplus K_{\lambda, \delta^n, \delta^n} S_{\lambda}(A \otimes B),$$

where $K_{\lambda, \delta^n, \delta^n}$ are Kronecker coefficients.

The alternate description is that

$$\mathbb{C}[GL(W).det_n] = Sym(\mathbb{C}^p \otimes \mathbb{C}^n \otimes \mathbb{C}^n)^{SL_n \times SL_n}$$

If λ has 2 parts (i.e. $p = 2$), then the coefficients $K_{2, \delta^n, \delta^n}$ are known. This is because the ring of invariants in this case is well known to be isomorphic to the ring of semi-invariants of a Kronecker quiver. see [45].

The connection is that $\mathbb{C}^p \otimes \mathbb{C}^q \otimes \mathbb{C}^r$ can be viewed as p copies of $\mathbb{C}^q \otimes \mathbb{C}^r$, i.e. p maps from \mathbb{C}^q to \mathbb{C}^r i.e. the representation space of a quiver with 2 vertices 1,2 and p arrows from vertex 1 to vertex 2, of dimension vector (q, r) . This is a generalized Kronecker quiver, for 2 arrows we get just the Kronecker quiver. If we look at occurrences of $S_{\lambda} \mathbb{C}^p \otimes S_{\mu} \mathbb{C}^q \otimes S_{\nu} \mathbb{C}^r$ in there, with λ having 2 parts, the same representation occurs with the same multiplicity in the coordinate ring, as for $p = 2$. In that case this is Kronecker quiver.

The Kronecker coefficients can be calculated using the Sylvester formula for the covariants of binary forms. Computing the Kronecker coefficients in general is a very difficult problem. For the computational complexity of these computations, see [7]. Immediately after the workshop, Manivel was able to make progress on this problem, [35].

There are special cases when, $K_{\lambda, \mu, \nu}$, are known to be computable in polynomial time. They are as follows. (1) one of partitions has one row. (2) two of the partitions have two rows. (3) two partitions are hooks. (4) one partition is a hook and another has two rows. (For these results and more, see [42–44]). We want to find conditions which will determine when $K_{\lambda, \delta^n, \delta^n}$ is zero or not. A central question is, “Is this decidability problem in P ?” Mumfley conjectures that this decidability problem is in P . However, this is an open question, [37]. Another possibly interesting article on this topic is [15].

Computational efforts have determined that $K_{\lambda, \delta^n, \delta^n}$ is almost never zero. In fact Derksen and Landsberg compared what occurred in the symmetric algebra in each degree. They did this in order to have a first approximation as to which modules could even be modules of polynomials. Unfortunately, not only did each module occur, but with high multiplicity.

We want to find conditions which will determine when $K_{\lambda, \delta^n, \delta^n}$ is zero or not for general p . Ultimately, the success of this line of reasoning will depend on asymptotic questions. It becomes necessary to describe a certain polytope which is asymptotically nonzero. There is a hope that this method will be reasonable because the $K_{\lambda, \delta^n, \delta^n}$ are not just generic Kronecker coefficients, they have a special format.

The idea was advanced to use the moment map to establish the vertices and/or faces of the cone

$$C(m, n, p) = \{(\lambda, \mu, \nu) \mid K_{\lambda, \mu, \nu} \neq 0\}$$

This is an interesting problem, with partial results established by Manivel during the workshop.

Step 2: The second example is the permanent $v = \text{perm}_m$. We need to consider

$$\mathbb{C}[GL(W).perm_m] = \bigoplus M_\lambda S_\lambda(A \otimes B),$$

where $M_\lambda = \Sigma_{\mu, \nu}$, $K_{\lambda, \mu, \nu}$, and we require that μ (resp. ν) is such that $[S_\mu A]_\circ^{\otimes n} \neq 0$ (resp. $[S_\nu B]_\circ^{\otimes n} \neq 0$).

This is seen to be a more tractable question, and there are a few papers on this. For $n = 3$ Weyman could show the formula for $[S_\mu A]_\circ$ as an S_3 -module.

Step 3: The final question is what happens when we expand to more variables, i.e. when making the transition from perm_m to $l^{n-m}\text{perm}_m$. Now there are n^2 variables and we lose the nice stability property.

Mulmuley and Sohoni tell us to consider partial stability, or (R, P) - stability. We must enlarge our group action from $GL(W)$, (where W is a m -dimensional subspace of V), to get a larger stabilizer of the form:

$$\begin{pmatrix} Mat_{n \times n} \mathbb{C} & * \\ 0 & * \end{pmatrix}$$

If we take the natural parabolic, $P \subset GL(W)$, we have a decomposition, $P = KU$, where U is a unipotent radical and K is the Levi factor. Note that $GL(W)$ is a reductive group. We choose a reductive subgroup $R \subset K$ which has the same rank (in the sense of the rank of the maximal torus). Then we choose $v \in PV$ such that the following properties hold:

- 1) v is R -stable (The orbit in affine space is closed under SL action.)
- 2) $U \subset G_v \subset P$

Then Mulmuley and Sohoni have a result which is a partial understanding of $\mathbb{C}[\overline{G.v}]$ in terms of $\mathbb{C}[\overline{R.v}]$. Because of the R -stability, they can gain some understanding from $\overline{R.v}$. This stability allows for a significant reduction: Instead of studying the $G = GL(n^2)$ -module structure of $\mathbb{C}[\overline{G.l^{n-m}\text{perm}_m}]$, we can study the $R = SL(m^2)$ -module structure of $\mathbb{C}[\overline{R.perm_m}]$, (See Theorem 6.2 [39] and the remarks following).

The addition of the linear factor l^{n-m} is also related to the Weyman-Kempf geometric technique of calculating syzygies, [53]. To this end one notices that the subvariety

$$Y = \{f \in S_d(W) \mid f = l^s g\}$$

where l is a linear form, can be treated by means of the geometric technique. One just has to consider the incidence variety

$$Z = \{(f, l) \in S_d(W) \times \mathbb{P}(W) \mid f = l^s g\},$$

which gives a desingularization of the orbit closure. Here we identify the projective space $\mathbb{P}(W)$ with the set of non-zero linear forms in W (modulo the usual equivalence relation). There are technical difficulties, as Y is usually not projectively normal.

4. OPEN PROBLEMS

- (1) In the case $n = 3$, compute the difference between $GL_{n^2}.det_n$ and $Mat_{n^2 \times n^2}.det_n$. One containment is obvious. In order to understand why the other containment could fail, consider the following example: The set $Mat_{2 \times 2}.(x^d + y^d)$ does not contain the GL_2 -orbit, $GL_2.(x^d + y^d)$ for $d \geq 4$. This example shows that one must be careful with orbits and closures as the behavior is not always as expected.

In general, compare $\overline{GL_{n^2}.det_n}$ to $Mat_{n^2 \times n^2}.det_n$. Note that while GL_{n^2} is a group, $Mat_{n^2 \times n^2}$ is an algebraic monoid.

- (2) (Landsberg) If the two orbits in (1) are not equal, is there a polynomial $p(n)$ such that $Mat_{p(n)^2 \times p(n)^2}.det_{p(n)}$ contains $\overline{l^{p(n)-n}GL_{n^2}.det_n}$ for some linear form l ?

The affirmative answer to this question would imply that $VP = \overline{VP}$.

- (3) Let $R = \text{Sym}(E \otimes F \otimes G) = \bigoplus_{\lambda, \mu, \nu} (S_\lambda E \otimes S_\mu F \otimes S_\nu G)^{K_{\lambda\mu\nu}}$, with $K_{\lambda\mu\nu}$ the Kronecker Coefficients. Let $\Delta(E, F, G) = \{(\lambda, \mu, \nu) \mid \exists n \text{ such that } S_{n\lambda} E \otimes S_{n\mu} F \otimes S_{n\nu} G \subset R\}$. It is known that $\Delta(E, F, G)$ is a rational polyhedral cone. **Determine the facets of this cone.**

Some cases of this question are already known. In the case of $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$, some inequalities on the facets are known. In [25], Franz considered the case when $\dim(E) = \dim(F) = \dim(G)$.

- (4) The case when $\dim(E) = 3$ and $\dim(F) = \dim(G) = n$. Compute the ring of invariants,

$$\text{Sym}(E \otimes F \otimes G)^{SL(F) \times SL(G)}$$

Note that if $\dim(E) = 2$, then this is the known Sylvester formula, which is related to the Kronecker coefficients. This problem would determine $K_{\lambda, \delta^n, \delta^n}$ for λ having ≤ 3 parts, (see [35]).

- (5) Consider $S_\lambda E$, where $|\lambda| = \sum \lambda_i$ is divisible by $\dim(E)$, (i.e. λ is in the root lattice). Let T be the maximal torus of $SL(E)$, i.e. the diagonal matrices with $\det = 1$. Let \mathcal{W} denote the Weyl group of $SL(E)$. Let $(S_\lambda E)_\circ$ be the zero weight space. Then $(S_\lambda E)_\circ = (S_\lambda E)^T$ is a representation of \mathcal{W} . For which λ does there exist a \mathcal{W} invariant? In other words, for which λ is $(S_\lambda E)_\circ^{\mathcal{W}} \neq 0$?

Remark 4.1. Questions 3 and 4 relate to the coordinate ring $\mathbb{C}[GL_{n^2}, \det_n]$, while questions 3 and 5 relate to the coordinate ring $\mathbb{C}[GL_{m^2}, \text{perm}_m]$. Question 3 leads to an asymptotic understanding, but it is a completely general question.

Remark 4.2. It is incredible how many important areas come to play.

Part 3. Matchgates and Holographic Algorithms

5. REPORT

In this group, the goal was to understand the work of J.-Y. Cai- P. Lu (see [8] for example) and L. Valiant (see [52]) in light of algebraic geometry and representation theory.

Let Γ be a planar graph. The FKT algorithm counts the number of perfect matchings for Γ in polynomial time. The goal of matchgates and holographic algorithms is to exploit this polynomial time algorithm to solve problems which were once unknown to be in P . The key to the polynomial time algorithm is that determinants and Pfaffians can be computed via linear algebra in polynomial time. The methods work when they can change a counting problem to one that just requires computing a determinant or Pfaffian.

An example of this is the following. Consider the problem $\#_7 Pl - NAE - ICE$ which asks the number (modulo 7) of orientations of a planar graph so that there are no sources and no sinks. It is possible to replace all of the vertices and edges of this graph with generators G and recognizers R and then try to model G and R with matchgates as follows. At each node, the signature $(0, 1, 1, 1, 1, 1, 0)$ is the truth table values required for the “not all equal” condition. At each edge, the signature $(0, 1, 1, 0)$ is truth table values required by the fact that each carries precisely one orientation. If these signatures can be realized by matchgates, then we will be able to compute $\#_7 Pl - NAE - ICE$ in polynomial time because this procedure changes the question about the old graph to a question about the new graph which can be done in polynomial time and gives the desired answer for the old graph.

For a given problem, what is desired is to know whether or not there exists a skew symmetric matrix associated to a particular signature. This can be determined by examining the so-called matchgate identities (MGI). If the desired signature satisfies MGI, then it is possible to compute that signature as the number of perfect matchings of a graph. If the desired signature does not satisfy MGI, all is not lost. Special basis changes are allowed as long as they are done in a way that is consistent with both the generator and the recognizer - more on this in a moment.

Let \mathbb{F} be a field. If Γ is a planar graph with γ nodes and \mathbb{F} -weighted edges, let $k(\Gamma)$ denote the skew symmetric matrix associated to Γ . If $I \subset [\gamma]$, let Pf_{2^I} denote the $\mathbb{C}^{2^{|I|}}$ -vector of principal sub-Pfaffians of $k(\Gamma)$ produced by omitting nodes indexed by $J \subset I$ and the adjacent edges. \underline{G} is a *standard signature* if it occurs Pf_{2^I} for some graph Γ (i. e. the MGI are satisfied). A signature G is realizable if there exists a linear map $f : \mathbb{C}^c \rightarrow \mathbb{C}^{2^b}$ so that $\underline{G} = f.G$. Note that this is just a linear map, and not even a group action. The map f is defined by taking a sufficient tensor power of a vector and implementing it as a matrix. Similarly,

a recognizer signature R is realizable if $R = \underline{R}.f$ for some standard signature \underline{R} and the same f as before. The goal is to find an f so that both R and G are realizable.

This problem has interesting connections to geometry that the participants hope to be able to exploit upon gaining further understanding of these connections. Consider $\text{Spec}((\mathbb{C}^{2^*})^k/MGI) = \mathbb{S}_k$. The MGI are well known to geometers, and the varieties that they cut out have been studied already. There is a canonical identification of the MGI variety, \mathbb{S}_k , with the spinor varieties, \mathcal{S}^{k-1} , given by the inclusion

$$\mathcal{S}_+^{k-1} \setminus \{0\} \sqcup \mathcal{S}_-^{k-1} \setminus \{0\} \hookrightarrow \mathbb{S}_k.$$

We also have maps $f : \mathbb{C}^c \rightarrow \mathbb{C}^{2^b}$ locally defined on the wires of the graph. So, given an explicit piece of data, $\mathbb{C}^{2^k} \otimes \mathbb{C}^{2^l}$, (assume $k < l$) and pairing information $[k] \hookrightarrow [l]$, and a map $f : \mathbb{C}^c \rightarrow \mathbb{C}^{2^k}$ as above, if $c = 2$ then the basis collapse theorem of Cai-Lu tells us that we can reduce the k to 2. It is unknown what happens if $c \neq 2$.

6. OPEN PROBLEMS

Exercise: Understand the space of signatures in the product of spinor varieties $\mathcal{S}_k \times \mathcal{S}_l$. What is the compatible action?

- (1) Using group actions, reprove the Cai-Lu Basis Collapse Theorem.

The vector space $\mathbb{C}^{2^k} \otimes \mathbb{C}^{2^l}$ has (at least) two useful descriptions: First, as the tensor product of spin representations: $\Delta_k \otimes \Delta_l$ - this reflects the action of the Spin groups on the space. (Recall the notation $\Delta_k = \bigwedge^{\text{even}} \mathbb{C}^N$.) The second description is as $(\mathbb{C}^2)^{\otimes k} \otimes (\mathbb{C}^2)^{\otimes l}$, and this reflects the action of the 2×2 matrices.

One may ask: Why not consider a larger space of actions, but the Cai-Lu result reduces to studying the action (actually just a mapping) of $\text{Mat}_2(\mathbb{C})$. (Gurvits comments that the spirit of this problem is like Quantum Computing and Unitary operations.)

The goal of this problem is that by rephrasing the proof in terms of geometry and representation theory, this will make the proof of the basis collapse theorem more transparent to geometers, and perhaps it will lead to a greater understanding. It is hoped that this greater understanding will lead to a proof of the next problem.

- (2) Study matchgates with colors instead of yes/no. Use the results of the previous problem to try to find a natural class of transformation for this setup. The idea is that one may be able to use a different group (GL_4 for example) in place of GL_2 . Does the Cai-Lu Theorem generalize? The expectation is that it will not, and in that case, this would be considered good news. This is related to a paper of Valiant which considered 3 colors and saw some success.

More specifically, if \underline{G} is a generator and \underline{R} is a recognizer, Cai-Lu considered the mapping by T , a $2^k \times 2$ matrix which sends the standard signatures $\underline{G}, \underline{R}$ to desirable non-standard signatures G, R . What if we consider the case that T is a $2^k \times 3$ matrix, or a $2^k \times 4$, etc, matrix?

- (3) Consider $(\mathbb{C}^2)^{\otimes k} \otimes ((\mathbb{C}^2)^*)^{\otimes l} \simeq \Delta_k \otimes \Delta_l$. We want to understand the diagonal GL_2 -action on points of \mathcal{S}_k as an orbit structure. Many cases are already known. For instance, Cai pointed out that there is a 2 dimensional space of matrices which leave \mathcal{S}_k invariant. What is desired is to understand and classify the points which have a 1 dimensional space of matrices which move the point off of the spinor variety. The reason is that if a point has this feature, there will then be 1 dimension's worth of freedom for the each factor in $\text{Seg}(\mathcal{S}_k \times \mathcal{S}_l)$. This in turn will tell which generators have a 1-parameter family of compatible recognizers.

Part 4. Entanglement and quantum information theory

7. REPORT

The contributions of G. Gour and D. Gross mention the long-standing additivity conjecture of quantum information theory. D. Gross posed a question which is closely related to this open problem – but has the

advantage of being independent of the measure used to quantify entanglement. In particular, there is no need to talk about non-algebraic quantities such as the “entropy of entanglement”.

The basic setting of the additivity conjecture is a linear space with a two-fold (horizontal and vertical) tensor product structure

$$(A_1 \otimes B_1) \\ \otimes (A_2 \otimes B_2).$$

Physically, the spaces $A = A_1 \otimes A_2$ and $B = B_1 \otimes B_2$ are associated with distinct observers, in a way that need not concern us. Let S_1 be any subspace of $A_1 \otimes B_1$ and likewise $S_2 \subset A_2 \otimes B_2$. Set $S = S_1 \otimes S_2$.

We are interested in the set of vectors in S which display the “least amount of entanglement” with respect to the partition A vs. B . The physical quantity “entropy of entanglement” measures the deviation of a vector $s \in S$ from being decomposable – of the form $s_A \otimes s_B$. (The precise definition will not be relevant for the question asked below). The additivity conjecture amounts to claiming that the set of least entangled vectors always contains one which is decomposable *with respect to the partition 1 versus 2*.¹ In other words, it is conjectured that if one aims to identify the global minimum of the entropy of entanglement in S , one may restrict attention to vectors of the form $s_1 \otimes s_2$, $s_i \in S_i$. It is likely that many researchers in the field would agree to slightly stronger and less precise statement that “generically” the global minimum is attained *only* on decomposable vectors.

D. Gross also asks the following consistency question. The optimization problem we set out to solve (find the least entangled vector in S) *does not refer to the vertical tensor product structure*. However, the conjectured statement does. So is the vertical decomposition intrinsic to the problem? Given just the data A, B and $S \subset A \otimes B$, when can one introduce such an additional tensor product structure and when is it unique?

If a negative answer could be given – i.e. if the tensor product structure is essentially uniquely determined by S – we would have learned an interesting fact about tensor product spaces. A positive answer would be an *extremely* interesting starting point for the search of a counter-example to the additivity conjecture.

Deciding additivity is considered the most high-profile open problem of quantum information. Any partial result would be met with great interest. There is no compelling evidence either in favor or against the conjecture.

The group working on entanglement and quantum information theory reports that while they did many calculations, they did not solve any of the open problems.

In addition, they computed for $s \in S_1 \otimes S_2$, the quantity $\min\{H(s)\}$, where H can be the entropy or the non-locality for example. They claim that it is also possible to compute

$$\min_{s_1 \in S_1} H(s_1) + \min_{s_2 \in S_2} H(s_2),$$

and in so doing, they find $s_1^{opt} \otimes s_2^{opt}$. It is unknown whether this is a local optimum. The following statement has been solved: If $s_1^{opt} \otimes s_2^{opt}$ is a local optimum, $s_1^{opt} \otimes s_2^{opt}$ is a critical point, and this implies that locally to first order that additivity works. Unfortunately, this tells nothing about the global problem.

8. OPEN PROBLEMS

1.) To which degree can one violate the Rank Multiplicativity (simpler version)?

Note: Very recently, the additivity conjecture for minimum output entropy was proven false by M.B. Hastings, [26]. So this problem is perhaps not so interesting as it was aimed at the additivity conjecture. None the less, we report on the activity from the workshop surrounding this problem.

Consider subspaces $S_1 \subset A_1 \otimes A_2$, $S_2 \subset B_1 \otimes B_2$, and let $r(S_i) = \min_{0 \neq s \in S_i} \{rank(s)\}$. Let $S = S_1 \otimes S_2$. We may consider $S \subset (A_1 \otimes A_2) \otimes (B_1 \otimes B_2)$, or under the natural braiding isomorphism, we can consider $S \subset (A_1 \otimes B_1) \otimes (A_2 \otimes B_2)$. It is known that there exist examples where $r(S) \neq r(S_1)r(S_2)$. Find examples

¹The conjecture pertains only to the case where the entropy of entanglement is used to measure the degree of non-decomposability of a vector. Analogous statements for several other measures – most notably the tensor rank – have recently been proven to be wrong.

of large violations. (Make it precise what it means to be “large”.) It should be noted that this is a toy example. There is not an automatic correspondence to the big problem for entropy or entanglement.

Following are two examples which were worked out during the workshop. Let $X \subset \text{Mat}_{n \times n}(\mathbb{R})$ be a linear subspace of real valued matrices, and let $Y = X \otimes X$. Define $\text{MinRank}(X)$ as a minimal rank of nonzero matrices in X . Clearly $(\text{MinRank}(X))^2 \geq \text{MinRank}(Y)$.

Proposition 8.1 (L. Gurvits). *The MinRank - multiplicativity conjecture, $(\text{MinRank}(X))^2 = \text{MinRank}(Y)$, is false over \mathbb{R} . Moreover, there can be an arbitrarily large decrement $(\text{MinRank}(X))^2 - \text{MinRank}(Y)$ over \mathbb{R} .*

Proof. Let I_k denote the $k \times k$ identity matrix. Consider the matrix, $M = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Let $X = \text{span}\{M, I_2\}$. Clearly, the minimum rank of any nonzero element in X is 2. The spectra $\sigma(M \otimes M) = (1, 1, -1, -1)$ and $M \otimes M$ is symmetric. Therefore, we find that $\text{rank}(M \otimes M + I_4) = \text{rank}(M \otimes M - I_4) = 2 < 4$. So MinRank multiplicativity is false.

For the arbitrary large decrement, modify the example as follows. Construct the linear space $S_{2n} = X \otimes I_n = \text{span}\{M \otimes I_n, I_2 \otimes I_n = I_{2n}\}$. Then the minimal rank of nonzero matrices in S_{2n} is $2n$, but $\text{rank}((M \otimes I_n) \otimes (M \otimes I_n) - I_{4n^2}) = 2n^2$ and this provides an arbitrarily large decrement. \square

Remark 8.2. S. Friedland observed that the same example shows that the entropy additivity conjecture is false over reals since $\log(2n) + \log(2n) > \log(2n^2)$.

- 2.) Let $\dim(S_i) = k_i$. Assume each S_i is generic in the sense that it is in general position in $A_i \otimes B_i$. Is it the case that you can have $r(S) = r(S_1)r(S_2)$? This is not just a dimension counting argument because $S_1 \otimes S_2$ may not be generic. In fact, it was commented that $S_1 \otimes S_2$ is never generic. (See Gurvits’s example for the real case above).
- 3.) Ambiguity of Tensor Decomposition. This area hasn’t been explored much.

Are there interesting situations which allow us to find a tensor product structure $A = A'_1 \otimes A'_2, B = B'_1 \otimes B'_2$, ($A'_i \neq A_i, B'_i \neq B_i$) and subspaces $S'_i \subset A'_i \otimes B'_i$ such that $S = S'_1 \otimes S'_2$? To qualify as “interesting”, S should contain no unentangled vector (of the form $s_A \otimes s_B$). Also, it would be desirable if the two decompositions $A = A_1 \otimes A_2$ and $A = A'_1 \otimes A'_2$ shared no decomposable vector. (There should be some more precise conditions given for a solution to be interesting).

The uniqueness of this decomposition is interesting. If you prove non-uniqueness, then you become a celebrity as it is related to an important problem in entanglement and quantum information theory.

REFERENCES

1. H. Abo, G. Ottaviani, and C. Peterson, *Induction for secant varieties of Segre varieties*, to appear in Trans. AMS, preprint math.AG/0607191.
2. L. Albera, A. Ferreol, P. Comon, and P. Chevalier, *Blind identification of overcomplete mixtures of sources (BIOME)*, Linear Algebra Appl. **391** (2004), 3–30.
3. J. Alexander and A. Hirschowitz, *La méthode d’Horace élatée: application à l’interpolation en degré quatre*, Invent. Math. **107** (1992), 585–602.
4. ———, *Polynomial interpolation in several variables*, J. Alg. Geom. **4** (1995), no. 2, 201–222.
5. ———, *Generic hypersurface singularities*, Proc. Indian Acad. Sci. Math. Sci. **107** (1997), no. 2, 139–154.
6. P. Bürgisser, *The complexity of factors of multivariate polynomials*, Foundations of Computational Mathematics **4** (2004), 369–396.
7. P. Bürgisser and C. Ikenmeyer, *The complexity of computing Kronecker coefficients*, Accepted for FPSAC 2008 <http://inst-mat.utalca.cl/fpsac2008>.
8. J. Cai and P. Lu, *Bases collapse in holographic algorithms*, IEEE Conference on Computational Complexity, IEEE Computer Society, 2007, pp. 292–304.
9. M. V. Catalisano, A. V. Geramita, and A. Gimigliano, *Secant varieties of $\mathbb{P}^1 \times \dots \times \mathbb{P}^1$ (n -times) are not defective for $n \geq 5$* , preprint: arXiv:0809.1701v1.
10. ———, *On the rank of tensors, via secant varieties and fat points. Zero-dimensional schemes and applications - Naples, 2000*, Queens Papers in Pure and Appl. Math. **123** (2002), 133–147.
11. ———, *Ranks of tensors, secant varieties of Segre varieties and fat points*, Linear Algebra Appl. **355** (2002), 263–285.
12. ———, *Erratum: Ranks of tensors, secant varieties of Segre varieties and fat points*, Linear Algebra Appl. **367** (2003), 247–348.

13. L. Chiantini and C. Ciliberto, *Weakly defective varieties*, Trans. Amer. Math. Soc. **354** (2002), no. 1, 151–178.
14. ———, *On the concept of k -sectant order of a variety*, J. London Math. Soc. **73** (2006), no. 2, 436–454.
15. M. Christandl, A. Harrow, and G. Mitchison, *On nonzero Kronecker coefficients and their consequences for spectra*, Commun. Math. Phys. **270** (2007), 575–585, arXiv:quant-ph/0511029v1.
16. P. Comon, *Canonical tensor decompositions*, Research Report RR-2004-17, I3S (17 June 2004).
17. ———, *Tensors decompositions. State of the art and applications*, IMA Conference of Mathematics in Signal Processing (December 18–20, 2000).
18. P. Comon, G. Golub, L-H. Lim, and B. Mourrain, *Symmetric tensors and symmetric tensor rank*, SIAM Journal on Matrix Analysis Appl. **30** (2008), no. 3, 1254–1279.
19. P. Comon and B. Mourrain, *Decomposition of quantics in sums of powers of linear forms*, Signal Processing **53** (1996), no. 2, 93–107, Special issue on High-Order Statistics.
20. P. Comon and M. Rajih, *Blind identification of under-determined mixtures based on the characteristic function*, Signal Processing **86** (2006), no. 9, 2271–2281, <http://dx.doi.org/10.1016/j.sigpro.2005.10.007>.
21. L. De Lathauwer and J. Castaing, *Blind identification of underdetermined mixtures by simultaneous matrix diagonalization*, Signal Processing, IEEE Transactions on **56** (2008), no. 3, 1096–1105.
22. L. De Lathauwer, J. Castaing, and J.-F. Cardoso, *Fourth-order cumulant-based blind identification of underdetermined mixtures*, Signal Processing, IEEE Transactions on **55** (2007), no. 6, 2965–2973.
23. G.R. Baron de Prony, *Essai expérimental et analytique: sur les lois de la dilatabilité de fluides élastique et sur celles de la force expansive de la vapeur del’alkool, à différentes températures*, Journal de l’École Polytechnique Floréal et Plairial **1** (1795), no. 22, 24–76.
24. V. de Silva and L.-H. Lim, *Tensor rank and the ill-posedness of the best low-rank approximation problem*, SIAM Journal on Matrix Analysis and Applications, Special Issue on Tensor Decompositions and Applications, to appear.
25. M. Franz, *Moment polytopes of projective \mathfrak{g} -varieties and tensor products of symmetric group representations*, Journal of Lie Theory **12** (2002), no. 2, 539–549.
26. M.B. Hastings, *A counterexample to additivity of minimum output entropy*, <http://arxiv.org/abs/0809.3972v2>.
27. J. Morton J.M. Landsberg, *The geometry of tensors: Applications to complexity, statistics and engineering*, in preparation.
28. V. Kac, *Some remarks on nilpotent orbits*, J. Algebra **61** (1980), no. 1, 189–918.
29. V. Kac and J. Dadok, *Polar representations*, J. Algebra **92** (1985), 504–524.
30. J.M. Landsberg and L. Manivel, *On the ideals of secant varieties of Segre varieties*, Found. of Comput. Math. (2004), 397–422.
31. Lieven De Lathauwer, *A link between the canonical decomposition in multilinear algebra and simultaneous matrix diagonalization*, SIAM Journal on Matrix Analysis and Applications **28** (2006), no. 3, 642–666.
32. Lieven De Lathauwer, Bart De Moor, and Joos Vandewalle, *Computation of the canonical decomposition by means of a simultaneous generalized schur decomposition*, SIAM Journal on Matrix Analysis and Applications **26** (2004), no. 2, 295–327.
33. A. Leahy, *A classification of multiplicity free representations*, J. Lie Theory **8** (1998), no. 2, 367–391.
34. Guillaume Malod and Natacha Portier, *Characterizing valiant’s algebraic complexity classes*, J. Complex. **24** (2008), no. 1, 16–38.
35. L. Manivel, *A note on certain Kronecker coefficients*, (2008), preprint: arXiv:0809.3710v1.
36. M. Mella, *Singularities of linear systems and the Waring problem*, Trans. Amer. Math. Soc. **358** (2006), no. 12, 5523–5538, MR 2238925 (2007h:14059).
37. K.D. Mulmuley, *Geometric complexity theory VI: The flip via saturated and positive integer programming in representation theory and algebraic geometry*, Technical Report TR-2007-04 (2007).
38. K.D. Mulmuley and M. Sohoni, *Geometric complexity theory I: An approach to the P vs. NP and related problems*, SIAM J. Comput. **31** (2001), no. 2, 496–526.
39. K.D. Mulmuley and M. Sohoni, *Geometric complexity theory, P vs. NP and explicit obstructions*, Advances in algebra and geometry (New Delhi), Hindustan Book Agency, 2003, pp. 239–261.
40. G. Ottaviani and C. Brambilla, *On partial polynomial interpolation*, preprint arXiv:0705.4448.
41. ———, *On the Alexander-Hirschowitz theorem*, J. Pure Appl. Algebra **212** (2008), 1229–1251.
42. J. B. Remmel, *A formula for the Kronecker products of Schur functions of hook shapes*, J. Algebra **120** (1989), no. 1, 100118.
43. ———, *Formulas for the expansion of the Kronecker products $S(m, n) \otimes S(1^{p-r}, r)$ and $S(1^k 2^l) \otimes S(1^{p-r}, r)$* , Discrete Math. **99** (1992), no. 1–3, 265–287.
44. M. H. Rosas, *The Kronecker product of Schur functions indexed by two-row shapes or hook shapes*, J. Algebraic Combin. **14** (2001), no. 2, 153–173.
45. A. Skowroński and J. Weyman, *The algebras of semi-invariants of quivers*, Transform. Groups **5** (2000), no. 4, 361–402.
46. V. Strassen, *Rank and optimal computation of generic tensors*, Linear Algebra Appl. **52/53** (1983), 645–685, MR 709378 (85b:15039).
47. A. Terracini, *Sulle vk per cui la varietà a degli sh $h + 1$ -secanti ha dimensione minore dell’ordinario*, Rend. Circ. Mat. Palermo **Selecta vol. I** (1911), 392–396.
48. S. Toda, *PP is as hard as the polynomial-time hierarchy*, SIAM J. Comput. **20** (1991), no. 5, 865–877.

49. ———, *Classes of arithmetic circuits capturing the complexity of computing the determinant*, IEICE TRANSACTIONS on Information and Systems **E75-D** (1992), no. 1, 116–124, Special Section PAPER (Special Section on Theoretical Foundations of Computing).
50. S. Toda and M. Ogiwara, *Counting classes are at least as hard as the polynomial-time hierarchy*, SIAM J. Comput. **21** (1992), no. 2, 316–328.
51. L. Valiant, *Completeness classes in algebra*, STOC '79: Proceedings of the eleventh annual ACM symposium on Theory of computing (New York, NY, USA), ACM, 1979, pp. 249–261.
52. ———, *Holographic algorithms*, SIAM J. Comput. **37** (2008), no. 5, 1565–1594.
53. J. Weyman, *Cohomology of Vector Bundles and Syzygies*, Cambridge Univ. Press, Cambridge, 2003, [Cambridge Tracts in Mathematics 149].

DEPT. OF MATHEMATICS, MAILSTOP 3368, TEXAS A &M UNIVERSITY, COLLEGE STATION, TX 77843-3368
E-mail address: oeding@math.tamu.edu