

A LIST OF OPEN PROBLEMS AND QUESTIONS

This document was produced as part of the activities during the Topology and Geometry of the Moduli space of Curves workshop in March 2005; it is a component of a larger resource site at <http://www.aimath.org/WWN/modspacecurves/>

Please send updates, additions, comments, corrections, etc. to either

- *Jeff Giansiracusa* (giansira@maths.ox.ac.uk)
- *or Daves Maulik* (dmaulik@math.princeton.edu).

This document originated as a rough transcription of the problems and questions generated during a discussion session towards the close of the workshop. Hopefully it will continue to evolve into a repository of new questions about the moduli space of curves. Towards this goal, readers are encouraged to submit new questions and information regarding developments on the questions listed here.

(Some of these problems may have already been solved, and the bibliographic data here is incomplete, so the reader is advised to do the usual background reading before investing time in these questions. Conversely, if you have any information on recent work or useful articles, please let us know so that we can keep this page current.)

(1) (Vakil, really Looijenga) Define:

- $a(\mathcal{M}_g) = g - 2$,
- $a(\mathcal{M}_{g,n}) = g - 1$ (for $n > 0$),
- $a(\mathcal{M}_{g,n}^c) = 2g - 3 - n$,
- $a(\mathcal{M}_{g,n}^{\text{rational tails}}) = g + n - 2$, and
- $a(\mathcal{M}_{g,n}^{\leq k \text{ rational components}}) = g - 1 + k$.

(see compact type for definitions of the partial compactifications referred to in the final three entries above.)

Question: Is it true that $\mathcal{M}_{g,n}^*$ has the homotopy type of a complex of (real) dimension $a(\mathcal{M}_{g,n}^*) + \dim_{\mathbb{C}} \mathcal{M}_{g,n}^*$? If not, is the cohomological dimension $\leq a(\mathcal{M}_{g,n}^*) + \dim_{\mathbb{C}} \mathcal{M}_{g,n}^*$? What about quasicohherent cohomological dimension $C \leq a(\mathcal{M}_{g,n}^*)$? Or with cohomology of ℓ -adic sheaves, is the cohomological dimension $a(\mathcal{M}_{g,n}^*) + \dim_{\mathbb{C}} \mathcal{M}_{g,n}^*$? (Note that some of these have already been answered by Looijenga.)

These questions come from a master conjecture:

- (a) $\mathcal{M}_{g,n}^*$ can be covered by fewer than $a(\mathcal{M}_{g,n}^*) + 1$ affine varieties.
- (b) (slightly weaker) It can be stratified by affine varieties of codimension $\leq a$.

Here 1a implies 1b, which implies everything else above.

See [HL97] for the source of this series of questions, and [RV04] for more discussion.

- (2) (Sullivan) Find an explicit cell decomposition of \mathcal{M}_g (no marked points) or $\overline{\mathcal{M}}_{g,n}$. Sullivan points out that there is no known cell decomposition of $\mathcal{M}_{g,0}$. Looijenga suspects one might be given by a good choice of Morse function, for instance, the minimum length of closed geodesics. Schmutz has done some work here. Sarnak has conjectured that $\det \Delta$ gives a Morse function on \mathcal{M}_g . (Is there a reason for Sarnak to believe this conjecture true?)
- (3) (Bodigheimer) A question related to both of the previous problems: What is the Lusternik-Schnirelman category of \mathcal{M}_g ? This is a number, bounded above by the dimension, and bounded below by the *cup-length* of the cohomology ring (the largest number of elements in positive degree which have a non-zero cup product. Can anyone compute the cup-length of the moduli space?

- (4) (Getzler et al.) Consider the Hochschild cohomology of the Fukaya category of a compact Kahler manifold X . (Roughly speaking, the Fukaya category has Lagrangian subspaces as objects and intersections as morphisms.)
- (a) Is this (naturally) isomorphic to quantum cohomology? The problem here is to make the *known* map mathematically rigorous. *What does this have to do with what we know about $\overline{\mathcal{M}}_{0,3}$?*
- (b) Is the cyclic homology of the Fukaya category of X isomorphic to the Gromow-Witten theory of X ?

Costello's theorems imply that these conjectures are plausible. Costello comments that the existence of the map should be a purely analytic question.

- (5) (Bertram) Are there any stable torsion classes in $AH^*(\mathcal{M}_{g,n})$? (Here AH denotes the image of Chow in cohomology.)

(Mondello) Is there a stable torsion class represented by some algebraic cycle which geometers "can see"? e.g. $\lambda_g \delta_0$ on $\overline{\mathcal{M}}_g$ satisfies $2\lambda_g \delta_0 = 0$. Is it zero? Faber says yes, over a field of characteristic 2. Teleman suggests that there are lots of natural torsion classes in K -theory (arising from the Grothendieck-Riemann-Roch theorem); how do they look in cohomology? (Compare to 12 below.)

- (6) (Madsen) Consider the Torelli group $I_{g,n} = \ker \mathcal{M}_{g,n} \rightarrow \text{Aut}(H_1(F_g), \omega) \cong Sp_{2g}\mathbb{Z}$. Let $g \rightarrow \infty$. Do the κ_{2i} classes restrict to 0 in $H^*(I_{g,n})$? Igusa thinks the answer is yes...

- (7) (Madsen) There is a potentially interesting connection between outer automorphisms and mapping class groups. Let F_n denote the free group on n generators and $\text{Aut}(F_n)$ its automorphism group. **Conjecture:** $\tilde{H}^*(B\text{Aut}(F_\infty); \mathbb{Q}) = 0$. Daniel Biss (of Chicago) has

suggested that the diagram

$$\begin{array}{ccc} B\Gamma_\infty^+ & \longrightarrow & BSp_\infty(\mathbb{Z})^+ \\ \downarrow & & \downarrow \\ BAut(F_\infty)^+ & \longrightarrow & BGL_\infty(\mathbb{Z})^+ \end{array}$$

becomes homotopy cartesian after localization at \mathbb{Q} . Borel's computation of $H^*(BSp_\infty(\mathbb{Z}); \mathbb{Q})$ and $H^*(BGL_\infty(\mathbb{Z}); \mathbb{Q})$ in conjunction with the Madsen-Weiss calculation of $H^*(B\Gamma_\infty, 1; \mathbb{Q})$ implies the conjecture. Perhaps it might be better to replace $BGL_\infty(\mathbb{Q})^+$ by Waldhausen's $A(*)$ and $BSp_\infty(\mathbb{Z})^+$ with a symplectic analogue $ASp(*)$. (Note: $A(pt)$ is rationally equivalent to $BGL(\mathbb{Z})^+$, and $ASp(pt)$ is rationally equivalent to $BSp(\mathbb{Z})^+$.) The resulting diagram might then be homotopy cartesian even before localizing at \mathbb{Q} . (We have seen cohomology classes in $BAut$, but they all vanish as rank goes to infinity.)

Some related thoughts: What is the relation with the restriction of κ_{2i} to $H^*(\mathcal{M}_{g,n}^c)$? Is $\mathcal{M}_{g,n}^c$ a $K(\pi, 1)$? Is it true that $\pi_1 \mathcal{M}_{g,n}^c$ is precisely the kernel of the Johnson homomorphism? (answer to this last question is yes, by Hain...?)

- (8) (Vakil) Regarding Faber's conjecture, explain the intersection number part (values of products of κ classes in $R^{g-2}\mathcal{M}_g$). This is known by Virasoro methods, but this should not be the "real reason." What is the real reason? This is related to Morita's conjecture announced at this conference [Mor05] (Conjecture 1, p. 4). For that matter, prove Morita's conjecture! Perhaps this is the "right" way to prove Faber's intersection number conjecture.

- (9) (Mondello) For $\mathcal{M}_{g,n}^{\leq k}$ rational components, is $R^{g-1+k} \cong \mathbb{Q}$? What is \mathcal{R}^{g-1+k} generated by? When should we expect a 1-dimensional socle, and what should we expect for $\mathcal{R}(\mathcal{M}_{g,n}^{\leq k})$?

- (10) (Igusa) Are there operations which relate the stable classes on $B\Gamma_\infty$? We have $H_{spec}^*(\mathbb{C}\mathbb{P}_{-1}^\infty) \cong \mathbb{Z}[c_1] \cdot u$, where u is the Thom class in degree -2 . How is this reflected at the level of infinite loop spaces? What are the stable maps $\mathbb{C}\mathbb{P}_{-1}^\infty \rightarrow \mathbb{C}\mathbb{P}_{-1}^\infty$?
- (11) (Baldwin) Has anyone computed the intersection cohomology of $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d)$? These can be arbitrarily singular, but this is what intersection cohomology is designed for. Is there a good notion of the tautological ring here? Perhaps the virtual fundamental class plays the usual role of the fundamental class.
- (12) (Faber) From Ekedahl and van der Geer, λ_g is 0 on \mathcal{A}_g rationally but not integrally. The order in integral cohomology has been computed up to a factor of two. What is it? (Compare to 5 above.)
- (13) (Bertram) When will Getzler's paper on $\overline{\mathcal{M}}_1$ appear (even just as a preprint)? Conjecture: $t \rightarrow \infty$. Getzler comments that he does not like how this question is phrased.

- (14) (Ellenberg) Consider Hurwitz space (genus g , degree d). Could the cohomology stabilize as $g \rightarrow \infty$ with d fixed? The reason behind this question is that point counting over finite fields gives exactly the behavior we would expect if we had Harer stability in degree 2.

So, could some sort of Harer stability hold for some sort of Hurwitz schemes? Motivation for this question comes from work on number fields/function fields done in the '80s by Darskovksy and Wright.

The general philosophy is this: suppose we have a nice sequence of varieties $\{X_n\}_{n \in \mathbb{N}}$, and

$$\lim_{n \rightarrow \infty} \frac{\# \text{ points on } X_n(\mathbb{F}_q)}{q^{\dim X_n}}$$

exists for all q . Is this because of some version of Harer stability at play here?

- (15) (Tseng) Same question for $C \rightarrow BG$, with G a finite group. Tillmann says "yes" for $G = S^1$. More precisely, consider

$$EDiff(F_{g,1}) \times_{\text{Diff}(F_{g,1})} \text{Map}^\partial(F_{g,1}, BG)$$

for G connected, such as S^1 . This stabilizes by gluing in tori and the induced map on homology is an isomorphism in some range. Is this related?

- (16) (Sullivan) Fix a curve C and look at all unbranched covers of it. This gives points in \mathcal{M}_g . Do these become uniformly dense for any C (with respect to the Teichmuller metric) in a universally defined region of \mathcal{M}_g , g large? This would imply that, given curves C_1, C_2 , one could find covers \tilde{C}_1, \tilde{C}_2 which are arbitrarily close in the moduli space, the Siegel-Ehrenpreis problem.

- (17) (Morita) There are many numerical invariants that can be associated to the moduli space.

- (a) One may compute the signature of the cohomology ring of $\overline{\mathcal{M}}_{g,n}$.
- (b) Since $\overline{\mathcal{M}}_{g,n}$ is a rational cohomology manifold, there are Thom's rational Pontrjagin classes, and one may compute the rational L -genus with respect to these.
- (c) Lastly, $\overline{\mathcal{M}}_{g,n}$ is an orbifold of a complex manifold, so there are orbifold Chern classes, and hence orbifold Pontrjagin classes. Thus one may talk about the orbifold L -genus.

What are these numbers? Do they agree? Probably not, but their disagreement would tell us interesting information about the types of singularities in the moduli space. The difference between the signature and the rational L -genus detects geometric singularities. The difference between the rational L -genus and the orbifold L -genus detects complex analytic singularities.

One may similarly ask questions about the signature of the tautological ring, and many other variations on this theme.

- (18) (Sullivan) This question regards the algebraic structure on the homology of the free loop space of a manifold. There are maps

$$H_*LM \xrightarrow{\Delta} H_*LM \otimes H_*LM$$

and

$$H_*LM \otimes H_*LM \xrightarrow{\mu} H_*LM$$

First the naive question: what is the algebraic structure here? The spectral sequence converging to H_*LM has E_2 term a tensor product of a Hopf algebra (coming from the base M) and a Frobenius algebra (coming from the fibre). But the differential does not respect these structures.

A perhaps better question is: can we illuminate the situation by reformulating in terms of the category of space over M ? We have

$$\begin{array}{ccccc} LM & \longrightarrow & LM \times_M LM & \longrightarrow & LM \times LM \\ \downarrow & & \downarrow & & \downarrow \\ M & \xrightarrow{\text{id}} & M & \xrightarrow{\Delta} & M \times M \end{array}$$

The left square corresponds to the Frobenius algebra part, and the right square corresponds to the Hopf algebra part. So, what is the full algebraic structure here?

REFERENCES

- [HL97] R. Hain and E. Looijenga, *Mapping class groups and moduli spaces of curves*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 97–142. MR [MR1492535](#) ([99a:14032](#))
- [Mor05] S. Morita, *Cohomological structure of the mapping class group and beyond*, preprint, 2005.
- [RV04] M. Roth and R. Vakil, *The affine stratification number and the moduli space of curves*, Algebraic structures and moduli spaces, CRM Proc. Lecture Notes, vol. 38, Amer. Math. Soc., Providence, RI, 2004, pp. 213–227. MR [MR2096147](#)