

MOTIVIC GALOIS GROUPS

BRUNO KAHN

1. TANNAKIAN CATEGORIES ([26], CF. [7])

K field of characteristic 0, \mathcal{A} rigid tensor K -linear *abelian* category, L extension of K .

Definition 1. An L -valued **fibre functor** is a tensor functor $\omega : \mathcal{A} \rightarrow \text{Vec}_L$ which is *faithful* and *exact*.

Definition 2. \mathcal{A} is

- **neutralised Tannakian** if one is given a K -valued fibre functor
- **neutral Tannakian** if $\exists K$ -valued fibre functor
- **Tannakian** if $\exists L$ -valued fibre functor for some L .

Example 1. G affine K -group scheme, $\mathcal{A} = \text{Rep}_K(G)$, $\omega : \mathcal{A} \rightarrow \text{Vec}_K$ the forgetful functor.

(\mathcal{A}, ω) neutralised Tannakian category: $G_K := \text{Aut}^\otimes(\omega)$ is (canonically) the K -points of an affine K -group scheme $G(\omega)$.

Theorem 1 (Grothendieck-Saavedra [26]). a) For (\mathcal{A}, ω) as in Example 1, $G(\mathcal{A}, \omega) = G$.
 b) In general ω enriches into a tensor equivalence of categories

$$\tilde{\omega} : \mathcal{A} \xrightarrow{\sim} \text{Rep}_K(G(\mathcal{A}, \omega)).$$

c) *Dictionary* (special case): \mathcal{A} semi-simple $\iff G$ proreductive.

When \mathcal{A} Tannakian but not neutralised, need replace $G(\mathcal{A}, \omega)$ by a *gerbe* (or a groupoid): Saavedra-Deligne [8].

Theorem 2 (Deligne [8]). \mathcal{A} rigid K -linear abelian. Equivalent conditions:

- \mathcal{A} is Tannakian
- $\forall M \in \mathcal{A}, \exists n > 0: \Lambda^n(M) = 0$.
- $\forall M \in \mathcal{A}, \dim_{\text{rigid}}(M) \in \mathbf{N}$.

2. ARE MOTIVES TANNAKIAN?

Ideally, would like $Mot_{\text{num}}(k, \mathbf{Q})$ Tannakian, fibre functors given by Weil cohomologies H .
Two problems:

$$\begin{array}{ccc}
 Mot_H & \xrightarrow{H} & Vec_K^* \\
 \downarrow & \nearrow \text{???} & \\
 Mot_{\text{num}} & &
 \end{array}$$

- $Mot_{\text{num}}(k, \mathbf{Q})$ is *never Tannakian* because $\dim_{\text{rigid}}(X) = \chi(X)$ may be negative (e.g. X curve of genus g : $\chi(X) = 2 - 2g$).

Second problem: matter of commutativity constraint – need modify it.

Yields Grothendieck's *standard conjectures* ([13], cf. [20]):

- (HN) $\sim_H = \sim_{\text{num}}$.
- (C) $\forall X$ the Künneth components of $H(\Delta_X)$ are algebraic.

Another conjecture (B) (skipped):

- (HN) \Rightarrow (B) \Rightarrow (C).
- (HN) \iff (B) in characteristic 0.

Theorem 3 (Lieberman-Kleiman [19]). *Conjecture (B) holds for abelian varieties.*

Theorem 4 (Katz-Messing [18]). *Conjecture (C) is true if k finite.*

Corollary 1 (Jannsen [14]). *If k finite, a suitable modification $\widetilde{\text{Mot}}_{\text{num}}(k, \mathbf{Q})$ is (abstractly) Tannakian.*

Apart from this, wide open!

Definition 3. When $\widetilde{\text{Mot}}_{\text{num}}(k, \mathbf{Q})$ exists, the gerbe that classifies it is called the [pure] **motivic Galois group** $GMot_k$. H Weil cohomology with coefficients K : fibre of $GMot_k$ at H is proreductive K -group $GMot_{H,k}$.

More generally, \mathcal{A} thick rigid subcategory of Mot_{num} , get an “induced” Galois group $GMot(\mathcal{A})$ of \mathcal{A} , quotient of the motivic Galois group. E.g. \mathcal{A} thick rigid subcategory generated by $h(X)$: get the **motivic Galois group of X** $GMot_{H,k}(X)$ (of finite type).

Examples 2.

- (1) $\mathcal{A} =$ Artin motives (generated by $h(\text{Spec } E)$, $[E : k] < \infty$): $GMot(\mathcal{A}) = G_k$.
- (2) $\mathcal{A} =$ pure Tate motives (generated by L or $h(\mathbf{P}^1)$): $GMot(\mathcal{A}) = \mathbb{G}_m$.
- (3) $\mathcal{A} =$ pure Artin-Tate motives (put these two together): $GMot(\mathcal{A}) = G_k \times \mathbb{G}_m$.
- (4) E elliptic curve over \mathbf{Q} , $H = H_{Betti}$.
 - E not CM $\Rightarrow GMot_{H, \mathbf{Q}}(E) = GL_2$.
 - E CM $\Rightarrow GMot_{H, \mathbf{Q}}(E) =$ torus in GL_2 or its normaliser.

Example 3. Suppose Conjecture (HN) true.

- **Characteristic 0:** Betti cohomology yields (several) \mathbf{Q} -valued fibre functors, as long as $\text{card}(k) \leq \text{card}(\mathbf{C})$: $\text{Mot}_{\text{num}}(k, \mathbf{Q})$ is neutral. Comparison isomorphisms \Rightarrow isomorphisms between various motivic Galois groups.
- **Characteristic p :** $k \supseteq \mathbf{F}_{p^2}$ finite $\Rightarrow \text{Mot}_{\text{num}}(k, \mathbf{Q})$ is *not neutral*: if $K \subseteq \mathbf{R}$ or $K \subseteq \mathbf{Q}_p$, no K -valued fibre functor (Serre: endomorphisms of a supersingular elliptic curve = quaternion \mathbf{Q} -algebra nonsplit by \mathbf{R}, \mathbf{Q}_p).

3. CONNECTION WITH HODGE AND TATE CONJECTURES

3.1. Tate conjecture.

k finitely generated, $G_k := \text{Gal}(\bar{k}/k)$, $H = H_l$ ($l \neq \text{char } k$): the \otimes -functor

$$H_l : \text{Mot}_H \rightarrow \text{Vec}_{\mathbf{Q}_l}^*$$

enriches into a \otimes -functor

$$\hat{H}_l : \text{Mot}_H \rightarrow \text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k)^*.$$

Tate conjecture $\iff \tilde{H}_l$ *fully faithful* (it is faithful by definition).

Proposition 1. *Tate conjecture \implies Conjecture (B).*

Hence under Tate conjecture, Conjecture (C) holds and can modify commutativity constraint:

$$\tilde{H}_l : \widetilde{\text{Mot}}_H \rightarrow \text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k).$$

$(\text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k), \text{forgetful functor})$ neutralised Tannakian \mathbf{Q}_l -category with fundamental group Γ_k : for $V \in \text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k)$, $\Gamma_k(V) = \text{Zariski closure of } G_k \text{ in } GL(V)$.

Proposition 2 (folklore, cf. [27], [17]). *Assume Tate conjecture. Equivalent conditions:*

- *Conjecture (HN);*
- $\text{Im } \tilde{H}_l \subseteq \text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k)_{ss}$ (full subcategory of semi-simple representations).

Under these conditions, Mot_{num} Tannakian, reduce to Γ_k^{ss} (for $\text{Rep}_{\mathbf{Q}_l}^{\text{cont}}(G_k)_{ss}$) proreductive and canonical epimorphism

$$\Gamma_k^{ss} \twoheadrightarrow GMot_{H_l, k}.$$

In particular, $\forall X$, $GMot_{H_l, k}(X) = \text{Zariski closure of } G_k \text{ in } GL(H_l(X))$.

Delicate question: essential image of \tilde{H}_l ? Conjectural answers for k finite (see below) and k number field (Fontaine-Mazur [11]).

3.2. Hodge conjecture.

$\sigma : k \hookrightarrow \mathbf{C}$, $H = H_\sigma$: this time enriches into \otimes -functor

$$\hat{H}_\sigma : \text{Mot}_{H_\sigma} \rightarrow PHS_{\mathbf{Q}}^*$$

(graded pure Hodge structures over \mathbf{Q}). Hodge conjecture $\iff \hat{H}_\sigma$ *fully faithful*.

Proposition 3. *Hodge conjecture* \implies *Conjecture (B)* \iff *Conjecture (HN)*.

Hence, under Hodge conjecture, get modified fully faithful tensor functor

$$\tilde{H}_\sigma : \widetilde{\text{Mot}}_{\text{num}} \rightarrow PHS_{\mathbf{Q}}.$$

Latter category semi-simple neutralised Tannakian (via forgetful functor). If extend scalars to \mathbf{R} , fundamental group = Hodge torus $S = R_{\mathbf{C}/\mathbf{R}}\mathbb{G}_m$. Over \mathbf{Q} it is the [Mumford-Tate group](#) *MT*: for $V \in PHS_{\mathbf{Q}}$, $MT(V) = \mathbf{Q}$ - Zariski closure of S in $GL(V)$.

Hodge conjecture $\iff \forall X, GMot_{k, H_\sigma}(X) = MT(X) \subseteq GL(H_\sigma(X))$.

Sometimes gives proof of Hodge conjecture (for powers of X , X abelian variety)!

4. UNCONDITIONAL MOTIVIC GALOIS GROUPS

Want an unconditional theory of motives (not assuming the unproven standard conjectures)

4.1. First approach (Deligne, André).

Both are in characteristic 0.

- **Deligne** [10]: replace motives by systems of compatible realisations: motives for **absolute Hodge cycles** (systems of cohomology classes corresponding to each other by comparison isomorphisms). Gives semi-simple Tannakian category.
Hodge conjecture \Rightarrow absolute Hodge cycles are algebraic so same category.
- **André** [3]: only adjoin to algebraic cycles the inverses of the Lefschetz operators: motives for **cycles**. Gives semi-simple Tannakian category.
Conjecture (B) \Rightarrow motivated cycles are algebraic so same category.
(Hodge conjecture \Rightarrow Conjecture (B) so cheaper approach!)

A abelian variety over number field:

Theorem 5 (Deligne [9]). *Every Hodge cycle on A is absolutely Hodge.*

Corollary 2. *Tate conjecture \Rightarrow Hodge conjecture on A .*

Better:

Theorem 6 (André [3]). *Every Hodge cycle on A is motivated.*

Corollary 3. *Conjecture (B) for abelian fibrations on curves \Rightarrow Hodge conjecture on A .*

Tannakian arguments:

Theorem 7 (Milne [23]). *Hodge conjecture for complex CM abelian varieties \Rightarrow Tate conjecture for all abelian varieties over a finite field.*

Theorem 8 (André [4]). *A abelian variety over a finite field: every Tate cycle is motivated.*

4.2. Second approach (André-K): tensor sections.

\mathcal{A} pseudo-abelian \mathbf{Q} -linear category, \mathcal{R} Kelly radical of \mathcal{A} (like Jacobson radical of rings): smallest ideal such that \mathcal{A}/\mathcal{R} semi-simple.

If \mathcal{A} tensor category, \mathcal{R} may or may not be stable under \otimes . True e.g. if \mathcal{A} Tannakian.

Theorem 9 (André-K [6]). *Suppose that \mathcal{R} is \otimes -ideal, $\mathcal{A}(\mathbf{1}, \mathbf{1}) = \mathbf{Q}$ and $\mathcal{R}(M, M)$ nilpotent ideal of $\mathcal{A}(M, M)$ for all M . Then the projection functor*

$$\mathcal{A} \rightarrow \mathcal{A}/\mathcal{R}$$

has tensor sections, and any two are tensor-conjugate.

Application:

H classical Weil cohomology,

$$\mathcal{A} = \text{Mot}_H^\pm(k, \mathbf{Q})$$

$$:= \{M \in \text{Mot}_H(k, \mathbf{Q}) \mid \text{sum of even Künneth projectors of } M \text{ algebraic}\}.$$

Then \mathcal{A} satisfies assumptions of Theorem 9: in characteristic 0 by comparison isomorphisms, in characteristic p by Weil conjectures.

Theorem 10 (André-K [5]). a) $\text{Mot}_{\text{num}}^\pm := \text{Im}(\text{Mot}_H^\pm \rightarrow \text{Mot}_{\text{num}})$ independent of H .

b) Can modify commutativity constraints in Mot_H^\pm and $\text{Mot}_{\text{num}}^\pm$, yielding $\widetilde{\text{Mot}}_H^\pm$ and $\widetilde{\text{Mot}}_{\text{num}}^\pm$.

c) Projection functor $\widetilde{\text{Mot}}_H^\pm \rightarrow \widetilde{\text{Mot}}_{\text{num}}^\pm$ has tensor sections σ ; any two are tensor-conjugate.

$$\begin{array}{ccc} \text{Mot}_H & \xrightarrow{H} & \text{Vec}_K^* \\ \downarrow & \nearrow \text{???} & \\ \text{Mot}_{\text{num}} & & \end{array}$$

$$\begin{array}{ccc} \widetilde{\text{Mot}}_H^\pm & \xrightarrow{H} & \text{Vec}_K \\ \downarrow \sigma & \nearrow H \circ \sigma & \\ \widetilde{\text{Mot}}_{\text{num}}^\pm & & \end{array}$$

Variant with

$$\text{Mot}_H^*(k, \mathbf{Q}) := \{M \in \text{Mot}_H(k, \mathbf{Q}) \mid \text{all Künneth projectors of } M \text{ algebraic}\}.$$

5. DESCRIPTION OF MOTIVIC GALOIS GROUPS

Assume all conjectures (standard, Hodge, Tate).

5.1. In general:

Short exact sequence

$$1 \rightarrow GMot_{\bar{k}} \rightarrow GMot_k \rightarrow G_k \rightarrow 1$$

Last morphism: G_k corresponds to motives of 0-dimensional varieties (Artin motives). The group $GMot_{\bar{k}}$ is connected, hence $= GMot_k^0$.

If $k \subseteq k'$, $GMot_{k'}^0 \twoheadrightarrow GMot_k^0$ (but not iso unless k'/k algebraic: otherwise, “more” elliptic curves over k' than over k).

Conjecture (C) \Rightarrow [weight grading](#) on Mot_{num} \iff central homomorphism

$$w : \mathbb{G}_m \rightarrow GMot_k.$$

On the other hand, Lefschetz motive gives homomorphism

$$t : GMot_k \rightarrow \mathbb{G}_m$$

and $t \circ w = 2$ (-2 with Grothendieck’s conventions).

5.2. Over a finite field:

Theorem 11 (cf. [22]). *a) Mot_{num} generated by Artin motives and motives of abelian varieties.*

b) Essential image of \tilde{H}_l : l -adic representations of G_k whose eigenvalues are Weil numbers.

Uses Honda's theorem [16]: every Weil orbit corresponds to an abelian variety.

Corollary 4. *$GMot_k^0 =$ group of multiplicative type determined by action of $G_{\mathbf{Q}}$ on Weil numbers.*

Even though $\widetilde{\text{Mot}}_{\text{num}}$ not neutral, $GMot_k^0$ abelian so situation not so bad!

5.3. Over a number field:

$S := (GMot_k^0)^{ab}$: the Serre protorus: describe its character group $X(S)$:

$$\mathbf{Q}^{cm} = \bigcup \{E \mid E \text{ CM number field}\}$$

Complex conjugation c central in $Gal(\mathbf{Q}^{cm}/\mathbf{Q})$ (largest Galois subfield of $\bar{\mathbf{Q}}$ with this property).

Definition 4. $f : Gal(\mathbf{Q}^{cm}/\mathbf{Q}) \rightarrow \mathbf{Z}$ **CM type** if $f(s) + f(cs)$ independent of s . $G_{\mathbf{Q}}$ acts on CM types by $\tau f(s) = f(\tau s)$.

Theorem 12 ([24]). $X(S) = \mathbf{Z}[CM \text{ types}]$.

Can also describe the centre C of $GMot_k^0$ (pro-isogenous to S), etc.: cf. [25].

6. MIXED (TATE) MOTIVES

Expect Tannakian category of **mixed motives**

$$\mathrm{Mot}_{\mathrm{num}}(k, \mathbf{Q}) \subset \mathrm{MMot}(k, \mathbf{Q})$$

with socle $\mathrm{Mot}_{\mathrm{num}}(k, \mathbf{Q})$, classifying non smooth projective varieties. Corresponding motivic Galois group extension of $GMot_k$ by a pro-unipotent group (or gerbe).

Constructions of MMot :

- Conjecturally, heart of “motivic t -structure” on DM (Deligne, Beilinson: cf. Hanamura [15]).
- In characteristic 0: explicit category constructed by Nori.
- Over a finite field: Tate conjecture $\Rightarrow \mathrm{Mot}_{\mathrm{num}} = \mathrm{MMot}$ (cf. [22]).
- Can settle for subcategory: mixed Tate motives TMMot_k . Exists unconditionally if k number field (cf. Levine’s talk and [21]).

Goncharov [12]: $\text{TMMot}_{\mathbf{Z}}$ (mixed Tate motives over \mathbf{Z}) defined as full subcategory of $\text{TMMot}_{\mathbf{Q}}$ by non-ramification conditions.

Γ the motivic Galois group corresponding to $\text{TMMot}_{\mathbf{Z}}$: Proreductive quotient of Γ is \mathbb{G}_m (see above).

Theorem 13 (Goncharov [12]). *Action of \mathbb{G}_m on prounipotent kernel U yields a grading on $\text{Lie}(U)$: for this grading, $\text{Lie}(U)$ is free with one generator in every odd degree ≤ -3 .*

- [1] Hodge cycles, motives and Shimura varieties, Lect. Notes in Math. **900**, Springer, 1982.
- [2] Motives, Proc. Symp. pure Math. **55** (I), AMS, 1994.
- [3] Y. André *Pour une théorie inconditionnelle des motifs*, Publ. Math. IHÉS **83** (1996), 5–49.
- [4] Y. André *Cycles de Tate et cycles motivés sur les variétés abéliennes en caractéristique p* , preprint, 2003.
- [5] Y. André, B. Kahn *Construction inconditionnelle de groupes de Galois motiviques*, C. R. Acad. Sci. Paris **334** (2002), 989–994.
- [6] Y. André, B. Kahn *Nilpotence, radicaux et structures monoïdales* (with an appendix of P. O’Sullivan), Rendiconti Sem. Math. Univ Padova **108** (2002), 107–291.
- [7] L. Breen *Tannakian categories*, in [2], 337–376.
- [8] P. Deligne *Catégories tannakiennes*, in The Grothendieck Festschrift, Progress in Math. **87** (II), Birkhäuser, 1990, 111–198.
- [9] P. Deligne *Hodge cycles on abelian varieties* (notes by J. S. Milne), in [1], 9–100.
- [10] P. Deligne, J.S. Milne *Tannakian categories*, in [1], 101–228.
- [11] J.-M. Fontaine, B. Mazur *Geometric Galois representations*, in Elliptic curves, modular forms, and Fermat’s last theorem (J. Coates, S.T. Yau, eds.), Intern. Press, Cambridge, MA, 1995, 41–78.
- [12] A. Goncharov *Multiple polylogarithms and mixed Tate motives*, preprint, 2001.
- [13] A. Grothendieck *Standard conjectures on algebraic cycles*, Algebraic Geometry — Bombay Colloquium, 1968, Oxford, 1969, 193–199.
- [14] U. Jannsen *Motives, numerical equivalence and semi-simplicity*, Invent. Math. **107** (1992), 447–452.
- [15] M. Hanamura *Mixed motives and algebraic cycles, III*, Math. Res. Lett. **6** (1999), 61–82.
- [16] T. Honda *Isogeny classes of abelian varieties over finite fields*, J. Math. Soc. Japan **20** (1968), 83–95.
- [17] B. Kahn *On the semi-simplicity of Galois actions*, to appear in the Rendiconti Sem. Mat. Univ. Padova **112** (2004).
- [18] N. Katz, W. Messing *Some consequences of the Riemann hypothesis for varieties over finite fields*, Invent. Math. **23** (1974), 73–77.
- [19] S. Kleiman *Algebraic cycles and the Weil conjectures*, in Dix exposés sur la cohomologie des schémas, North-Holland, Amsterdam, 1968, 359–386.
- [20] S. Kleiman *The standard conjectures*, in [2], 3–20.
- [21] M. Levine *Tate motives and the vanishing conjectures for algebraic K-theory*, in Algebraic K-theory and algebraic topology (Lake Louise, 1991), NATO ASI Series, Ser. C **407**, Kluwer, 1993, 167–188.
- [22] J. S. Milne *Motives over finite fields*, in [2], 401–459.
- [23] J. S. Milne *Lefschetz motives and the Tate conjecture*, Compos. Math. **117** (1999), 45–76.
- [24] J. S. Milne, K-Y. Shih *Langland’s construction of the Taniyama group*, in [1], 229–260.
- [25] J.-P. Serre *Propriétés conjecturales des groupes de Galois motiviques*, in [2], 377–400.
- [26] N. Saavedra Rivano *Catégories tannakiennes*, Lect. Notes in Math. **265**, Springer, 1972.
- [27] J. T. Tate *Conjectures on algebraic cycles in l -adic cohomology*, in [2], 71–83.