

Mean Values of L-Functions and Symmetry

J. B. Conrey and D. W. Farmer

1 Introduction

Katz and Sarnak [KS1] have introduced the idea of a family of L-functions with an associated symmetry type. The symmetry type has been shown to govern the distribution and spacing of zeros in the function field case [KS2], and there is strong numerical evidence [Ru] that it governs the behavior of zeros in more general cases. The paper [ILS] also shows clear evidence that the symmetry type governs the distribution of low-lying zeros of a wide variety of L-functions. In this paper we give evidence that the symmetry type of a family of L-functions governs the behavior of mean values of the L-functions.

The best-understood mean values are the $2k$ th moments of the Riemann ζ -function. We describe that case in detail, and then we compare this to the examples provided by the families of Katz and Sarnak.

It is a folklore conjecture that for every $k \geq 0$ there is a constant c_k such that

$$I_k(T) := \frac{1}{T} \int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^{2k} dt \sim c_k \log^{k^2} T$$

as $T \rightarrow \infty$. Classically, Hardy and Littlewood [HL] proved that

$$I_1(T) \sim \log T,$$

and Ingham [I] showed that

$$I_2(T) \sim \frac{1}{2\pi^2} \log^4 T,$$

Received 17 April 2000.

Communicated by Peter Sarnak.

so $c_0 = c_1 = 1$ and $c_2 = 1/2\pi^2$. No other mean values of the ζ -function have been established.

More information can be given about c_k , for Conrey and Ghosh [CG1] showed (assuming the Riemann hypothesis) that for all $k \geq 0$,

$$I_k \geq (1 + o(1)) \frac{\alpha_k}{\Gamma(1 + k^2)} \log^{k^2} T,$$

where α_k is an arithmetic constant given by

$$\alpha_k = \prod_p \left(1 - \frac{1}{p}\right)^{k^2} \sum_{j=0}^{\infty} \frac{d_k(p^j)^2}{p^j}. \quad (1.1)$$

Here $d_k(n)$ is the n th coefficient in the Dirichlet series for $\zeta(s)^k$; it is multiplicative and is given by

$$d_k(p^j) = \frac{\Gamma(k + j)}{\Gamma(k)j!}.$$

Gonek [G] extended the above result to include all $k > -1/2$.

Conrey and Ghosh [CG2] established properties of α_k . These included that α_k , regarded as a function of k , is entire of order 2, satisfies a symmetry $\alpha_k = \alpha_{1-k}$, and has all of its zeros on the line $\Re(k) = 1/2$. In addition, all of the zeros of the derivative da_k/dk have real parts equal to $1/2$ with two exceptions $a'_0 = a'_1 = 0$. Conrey and Gonek [CGo] give an asymptotic formula for $\log \alpha_k$ as $k \rightarrow \infty$. This formula is relevant to understanding the extremely large values of $\zeta(s)$.

Conrey and Ghosh [CG2] defined the number g_k implicitly by the (conjectural) formula

$$I_k(T) \sim g_k \frac{\alpha_k}{\Gamma(1 + k^2)} \log^{k^2} T. \quad (1.2)$$

The quantity g_k is natural in the following sense. All approaches to mean value theorems for Dirichlet series have relied on techniques from the theory of Dirichlet polynomials. The main tool there is the mean value theorem of Montgomery and Vaughan [MV]:

$$\int_0^T \left| \sum_{n=1}^N a_n n^{it} \right|^2 dt = \sum_{n=1}^N (T + O(n)) |a_n|^2.$$

To use this to obtain asymptotic formulae it is usually necessary to have $N \ll T$. Thus, it

is natural in some sense to measure the mean square of $\zeta(s)^k$ against the mean value of a Dirichlet polynomial approximation to $\zeta(s)^k$ using a polynomial of length T . It is easy to show that

$$\frac{1}{T} \int_0^T \left| \sum_{n \leq T} \frac{d_k(n)}{n^{1/2+it}} \right|^2 dt \sim \frac{a_k}{\Gamma(1+k^2)} \log^{k^2} T.$$

Thus, g_k is a measure of “how many polynomials of length T are needed to capture the mean square of $\zeta(s)^k$.”

The classical results of Hardy and Ingham can be phrased as $g_1 = 1$ and $g_2 = 2$. Recently, Conrey and Ghosh [CG3] made the conjecture that $g_3 = 42$. Still more recently, Conrey and Gonek [CGo] conjectured that $g_4 = 2024$. The conjectures for g_3 and g_4 are based on Dirichlet polynomial techniques. With current methods it is probably not possible to use those techniques to conjecture g_k for larger values of k . We return to the function g_k after giving a general discussion of mean values and a description of the situation for the families of L-functions of Katz and Sarnak.

We now describe the situation for some of the families of Katz and Sarnak. The mean values are of the shape

$$\frac{1}{Q^*} \sum_{\substack{f \in \mathcal{F} \\ c(f) \leq Q}} V\left(L_f\left(\frac{1}{2}\right)\right)^k \sim g_k \frac{a_k}{\Gamma(1+B(k))} (\log Q^A)^{B(k)}. \tag{1.3}$$

The L-functions are normalized to have a functional equation $s \leftrightarrow 1 - s$, so $L_f(1/2)$ is the “critical value.” Here we think of the family \mathcal{F} as being partially ordered by “conductor” $c(f)$, with Q^* the number of elements with $c(f) \leq Q$. We set $V(z) = z$ or $|z|^2$ depending on the symmetry type of the family. The parameters g_k and $B(k)$ depend only on the symmetry type and are integral for integral k , with $g_1 = 1$. The parameter a_k depends on the family in a natural way and is similar to the case of the ζ -function. The parameter A depends both on the symmetry type and the functional equation satisfied by the elements in the family (specifically, it depends on the degree of the functional equation in the relevant parameter). Examples are given below.

One of the symmetry types described by Katz and Sarnak is denoted O , for *orthogonal*. Examples of families with this symmetry type conjecturally include

- (a) the L-functions $L_f(s)$ associated with cusp forms $f \in S_m(\Gamma_0(1))$ of weight m for the full modular group,
- (b) the L-functions $L_f(s)$ with $f \in S_2(\Gamma_0(N))$ of weight 2 for the Hecke congruence group $\Gamma_0(N)$,

- (c) the twisted L-functions $L(s, \text{sym}^{2\ell+1}(f) \otimes \chi_d)$, where f is a self-dual cuspidal automorphic form on GL_2 and χ_d is a quadratic Dirichlet character mod $|d|$, provided that the Fourier coefficients of f have a Sato-Tate distribution, and
- (d) the twisted L-functions $L_f(s, \chi_d)$, where f is a self-dual cuspidal automorphic form on GL_m for some m , provided that the symmetric square L-function of f does not have a pole at $s = 1$.

To illustrate how the symmetry type is related to the mean values of the L-functions in the family, we note the following conjectures:

$$\frac{1}{m^*} \sum_{f \in S_m(\Gamma_0(1))} L_f\left(\frac{1}{2}\right)^k \sim g_k \frac{a_k}{\Gamma\left(1 + \frac{1}{2}k(k-1)\right)} (\log m^{1/2})^{(1/2)k(k-1)},$$

where m^* is the cardinality of S_m ,

$$\frac{1}{N^*} \sum_{f \in S_2(\Gamma_0(N))} L_f\left(\frac{1}{2}\right)^k \sim g_k \frac{a_k}{\Gamma\left(1 + \frac{1}{2}k(k-1)\right)} (\log N^{1/2})^{(1/2)k(k-1)},$$

where N^* is the cardinality of $S_2(\Gamma_0(N))$, and

$$\frac{1}{D^*} \sum_{|d| \leq D} L\left(\frac{1}{2}, \text{sym}^{2\ell+1}(f) \otimes \chi_d\right)^k \sim g_k \frac{a_k}{\Gamma\left(1 + \frac{1}{2}k(k-1)\right)} (\log D^{\ell+1})^{(1/2)k(k-1)},$$

where D^* is the number of quadratic characters with conductor not exceeding D .

Note that these families can be further broken into the even and odd forms, each of which is approximately half of the family. The average over the odd forms is identically zero because the associated L-functions vanish at the center of the critical strip.

We were unable to locate a reference for these conjectures. The above formulas were found, for example, using the Petersson formula and are based on results found in [D], [DFI], and [KMV]. From those papers we obtain $g_1 = 1$, $g_2 = 2$, $g_3 = 2^3$, and $g_4 = 2^7$.

In the notation of (1.3), the families with symmetry type O have $V(z) = z$ and $B(k) = (1/2)k(k-1)$. For the remaining parameter we have $A = \mathcal{A}$, where \mathcal{A} is the degree to which the parameter Q occurs in the functional equation. For example, the first two families above satisfy a functional equation of the form

$$\Phi(s) = \left(\frac{N^{1/2}}{2\pi}\right)^s \Gamma\left(s + \frac{m-1}{2}\right) L_f(s) = \varepsilon \bar{\Phi}(1-s).$$

That functional equation has degree 1/2 in both N and m aspect, so in the corresponding mean value we have $\Lambda = 1/2$.

Another symmetry type considered by Katz and Sarnak is denoted Sp, for *symplectic*. Examples conjecturally are

- (e) Dirichlet L-functions $L(s, \chi_d)$, where χ_d is a quadratic Dirichlet character mod $|d|$,
- (f) the symmetric square L-functions $L(s, \text{sym}^2(f))$ associated with $f \in S_m(\Gamma_0(1))$,
- (g) the twisted L-functions $L(s, \text{sym}^{2\ell}(f) \otimes \chi_d)$, where f is a self-dual cuspidal automorphic form on GL_2 , provided that the Fourier coefficients of f have a Sato-Tate distribution, and
- (h) the twisted L-functions $L_f(s, \chi_d)$, where f is a self-dual cuspidal automorphic form on GL_m for some m , provided the symmetric square L-function of f has a pole at $s = 1$.

The conjectured mean value in case (e) is

$$\frac{1}{D^*} \sum_{|d| \leq D} L\left(\frac{1}{2}, \chi_d\right)^k \sim g_k \frac{a_k}{\Gamma\left(1 + \frac{1}{2}k(k+1)\right)} (\log D^{1/2})^{(1/2)k(k+1)},$$

where D^* is the number of quadratic characters with conductor not exceeding D . This conjecture is based on work of Jutila [J] and Soundararajan [S]. In this case we have $g_1 = 1$, $g_2 = 2$, $g_3 = 2^4$, and it is conjectured that $g_4 = 3 \cdot 2^8$.

In case (f) we have the conjecture

$$\frac{1}{m^*} \sum_{f \in S_m(\Gamma_0(1))} L\left(\frac{1}{2}, \text{sym}^2(f)\right)^k \sim g_k \frac{a_k}{\Gamma\left(1 + \frac{1}{2}k(k+1)\right)} (\log m^{1/2})^{(1/2)k(k+1)},$$

where m^* is the cardinality of S_m . The cases (g) and (h) look just like (e).

In the notation of (1.3), families with symmetry type Sp have $V(z) = z$ and $B(k) = (1/2)k(k+1)$. The parameter Λ for the Sp families is determined in exactly the same way as for the O families.

A third symmetry type described by Katz and Sarnak is denoted U, for *unitary*. An example of a family with this symmetry type is

- (i) $L(s, \chi)$ for χ a character mod q .

The conjectured mean value in this case is

$$\frac{1}{Q^*} \sum_{|q| \leq Q} \sum_{\chi \bmod q} \left| L\left(\frac{1}{2}, \chi\right) \right|^{2k} \sim g_k \frac{a_k}{\Gamma(1+k^2)} (\log Q)^{k^2},$$

where the inner sum is over characters mod q , and Q^* is the number of characters with conductor at most Q .

For families with symmetry type U we have $V(z) = |z|^2$ and $B(k) = k^2$. For the remaining parameter we have $A = 2\mathcal{A}$, where \mathcal{A} is the degree to which the parameter Q appears in the functional equation. As example (i) suggests, we can think of the Riemann ζ -function as forming its own unitary family, where the mean values of the ζ -function correspond to averages of special values of the family $\{\zeta(1/2 + it)\}_{t \in \mathbb{R}}$.

We end this discussion by giving an example of the parameter α_k in the above formulas. The following is the α_k associated to with family of real Dirichlet L -functions $L(s, \chi_d)$, which has symmetry type Sp :

$$\alpha_k = \prod_p \frac{\left(1 - \frac{1}{p}\right)^{k(k+1)/2}}{\left(1 + \frac{1}{p}\right)} \left(\frac{1}{2} \left(\left(1 + \frac{1}{\sqrt{p}}\right)^{-k} + \left(1 - \frac{1}{\sqrt{p}}\right)^{-k} \right) + \frac{1}{p}\right).$$

While all of the above mean values are conjectural (except for certain small values of k), the various parameters in the formulas are fairly well understood, except for the constant g_k . Recently, Keating and Snaith [KeSn] have used techniques from random matrix theory to obtain conjectures for g_k for the symmetry types given above. This is described in Section 2.

In this paper we study the functions g_k in detail. We also report on calculations of the mean square of the L -function times a “mollifier” for each of the symmetry types described above. The results on g_k are summarized in the next section, and the results on mollifiers are presented in the following section. The remainder of the paper is devoted to results related to the various properties of the g_k functions.

2 Statement of results

The constants g_k occurring in the conjectured mean values of these L -functions is the most mysterious aspect of these mean values. Conjectures for g_k have recently been given, which we now describe.

The methods used to obtain conjectures for g_k are based on random matrix theory. This approach to the study of L -functions began with Montgomery’s work [M] on the pair correlation of zeros of the Riemann ζ -function, and it has been fruitful for establishing both rigorous and conjectural results about L -functions. The underlying idea is the conjecture that the zeros of L -functions are distributed on the critical line like the eigenvalues of matrices from the Gaussian unitary ensemble of large random Hermitian matrices. This is referred to as the “GUE” conjecture. This conjecture has been

corroborated, to a large extent, by the extensive computations of Odlyzko [O]. Heuristic explanations for the GUE conjecture have been given by Bogomolny and Keating [BK]. Now Keating and Snaith [KeSn] have announced conjectures for g_k that have been obtained by techniques from this theory.

Keating and Snaith’s conjecture in the unitary case is that

$$g_{\lambda,u} = \Gamma(1 + B_u(\lambda)) \lim_{N \rightarrow \infty} N^{-B_u(\lambda)} \prod_{j=1}^N \frac{\Gamma(j)\Gamma(j + 2\lambda)}{\Gamma(j + \lambda)^2}, \tag{2.1}$$

where $B_u(\lambda) = \lambda^2$. The conjecture in the orthogonal case is

$$g_{\lambda,o} = \Gamma(1 + B_o(\lambda)) \lim_{N \rightarrow \infty} N^{-B_o(\lambda)} 2^{2N\lambda-1} \prod_{j=1}^N \frac{\Gamma(N + j - 1)\Gamma\left(j - \frac{1}{2} + \lambda\right)}{\Gamma(N + j - 1 + \lambda)\Gamma\left(j - \frac{1}{2}\right)}, \tag{2.2}$$

where $B_o(\lambda) = (1/2)\lambda(\lambda - 1)$. And the conjecture in the symplectic case is

$$g_{\lambda,sp} = \Gamma(1 + B_{sp}(\lambda)) \lim_{N \rightarrow \infty} N^{-B_{sp}(\lambda)} 2^{2N\lambda} \prod_{j=1}^N \frac{\Gamma(N + j + 1)\Gamma\left(j + \frac{1}{2} + \lambda\right)}{\Gamma(N + j + 1 + \lambda)\Gamma\left(j + \frac{1}{2}\right)}, \tag{2.3}$$

with $B_{sp}(\lambda) = (1/2)\lambda(\lambda + 1)$.

We have recently learned that Brézin and Hikami [BH] have independently obtained the above conjectures in the Sp and O cases.

It is of key importance that formulas (2.1)–(2.3) agree with the known and conjectural values of g_k given earlier. These formulas for g_λ suggest that it would be natural to absorb the factor $\Gamma(1 + B(\lambda))$ into the definition of g_λ . However, doing so would obscure the arithmetic origins of the parameters in the mean values, and we would lose the significant fact that g_k is integral for k a positive integer.

In this paper, we develop the properties of the Keating-Snaith constants g_λ . We show that $g_\lambda/\Gamma(1 + B(\lambda))$ is a nonvanishing meromorphic function of order 2. We give a complete description of its pole locations, and we express $g_\lambda/\Gamma(1 + B(\lambda))$ in terms of the Barnes-Vignéras double Γ -function (see [V], [Sa], and [UN]).

Moreover, g_k is an integer for positive integer k , and we establish asymptotic formulas for g_k as $k \rightarrow \infty$. We remark that there are interesting patterns in the prime factorization of g_k . The exponents of small primes dividing g_k are quite irregular, and

in fact $p \nmid g_{k,U}$ for $k < p < k + \sqrt{p}$. For example,

$$\begin{aligned}
 g_{100,U} = & 2^{95} \cdot 3^{65} \cdot 5^{24} \cdot 7^{33} \cdot 11^{10} \cdot 13^{33} \cdot 17^{36} \cdot 19^{29} \cdot 23^{20} \cdot 29^{16} \cdot 31^{11} \cdot 37^{10} \cdot 41^{12} \\
 & \cdot 43^9 \cdot 47^4 \cdot 53^3 \cdot 59^7 \cdot 61^9 \cdot 67^{18} \cdot 71^{12} \cdot 73^{10} \cdot 79^6 \cdot 83^4 \cdot 89^2 \cdot 97 \cdot 113 \\
 & \cdot 127^5 \cdot 131^7 \cdot 137^9 \cdot 139^{10} \cdot 149^{16} \cdot 151^{17} \cdot 157^{20} \cdot 163^{24} \cdot 167^{26} \cdot 173^{30} \\
 & \cdot 179^{34} \cdot 181^{36} \cdot 191^{43} \cdot 193^{44} \cdot 197^{47} \cdot 199^{47} \cdot 211^{47} \cdot 223^{44} \cdot \dots \cdot 9973.
 \end{aligned}$$

Even more interesting are the patterns in the exponent of p in the prime factorization of g_k as $k \rightarrow \infty$. We show that there are continuous self-similar functions $c_p(x)$ such that if $k_j = [p^j x]$, then $v_p(g_{k_j,U}) \sim k_j c_p(x)$, and $v_p(g_{k_j,O}) \sim v_p(g_{k_j,Sp}) \sim (1/2)k_j c_p(x)$, where $v_p(n)$ is the power of p in the factorization of n . A graph of $c_3(x)$ is shown in Figure 1.




Figure 1

Note that these functions satisfy $c_p(x) = c_p(px)$. We establish other self-similarity properties of $c_p(x)$, and we also give an elegant formula for $c_p(x)$.

As a matter of interest, we also report that the Keating-Snaith constant $g_{1/2,U}$ can be used to conjecture the mean 1st moment of the ζ -function, and a calculation finds

$$g_{1/2,U} = \frac{\Gamma\left(\frac{5}{4}\right)\pi^{1/4}}{2^{1/6}} \exp\left(\frac{1}{4}\left(\frac{\zeta'}{\zeta}(2) - \gamma + 1\right)\right) \approx 1.0362329154\dots \tag{2.4}$$

It has been shown in [CG1] and [HB] that $g_{1/2,U}$, if it exists, satisfies $1 \leq g_{1/2,U} \leq 16/15$, assuming the Riemann hypothesis.

The paper is organized as follows. In Section 3 we discuss the mean square of L-functions times a mollifier. The remainder of the paper is devoted to establishing properties of g_λ as defined in (2.1)–(2.3). We write g_λ and $B(\lambda)$ to mean $g_{\lambda,X}$ and $B_X(\lambda)$ for X any one of U, O , or Sp . Results that are particular to one of the $g_{\lambda,X}$ are specifically

stated as such. In Section 4 we determine analytic properties of g_λ as a function of the complex number λ . This permits us to express g_λ in terms of the double Γ -function. In Section 5 we prove integrality properties of g_k for k a positive integer. In Section 6 we discuss the self-similar functions $c_p(x)$. In Section 7 we give an asymptotic formula for g_k as $k \rightarrow \infty$.

3 Mollified mean squares

We model our discussion on the mollified mean square required for Levinson’s method (see [Lev] and [C]). If

$$L(s)^{-1} = \sum_{n=1}^{\infty} \frac{m(n)}{n^s},$$

then our mollifier for $L(s)$ will be

$$M(s, f) = \sum_{n \leq y} \frac{m(n)}{n^s} f\left(\frac{\log \frac{y}{n}}{\log y}\right),$$

where $f(x)$ is a real polynomial with $f(0) = 0$, and $y = Y^\theta$ for some $\theta > 0$, with Y chosen appropriately for each family. We are interested in an asymptotic formula for the mean square of $L(s)M(s, f)$ for L in the families described in a previous section.

We first consider the unitary family, for which our model is the Riemann ζ -function. Let

$$\mathcal{M}_U(P, Q, \theta) := \frac{1}{T} \int_1^T \left| Q\left(\frac{-1}{\log T} \frac{d}{da}\right) \zeta\left(\frac{1}{2} + a + it\right) M\left(\frac{1}{2} + it, P\right) \right|_{a=0}^2 dt.$$

The length of the mollifier is T^θ . The following formula holds (see [C]) for $\theta < 4/7$:

$$\mathcal{M}_U(P, Q, \theta) \sim P(1)^2 Q(0)^2 + \frac{1}{\theta} \int_0^1 \int_0^1 \left(P'(x)Q(y) + \theta P(x)Q'(y) \right)^2 dx dy.$$

An interesting simple case is $\mathcal{M}_U(x, 1, \theta) \sim 1 + \theta^{-1}$.

For the orthogonal family, the mollifier has length \sqrt{X}^θ . The mollified mean value is

$$\mathcal{M}_O(P, Q, \theta) := \frac{1}{X^*} \sum_{c(f) \leq X} \left(Q\left(\frac{2}{\log X} \frac{d}{da}\right) \Phi_f\left(\frac{1}{2} + a\right) M\left(\frac{1}{2}, P\right) \right) \Big|_{a=0}^2.$$

For examples of applications of related mollified mean values, see [IS], [KM], and [KMOV]. If Q is even or odd, then for $\theta < 1$ we have

$$\mathcal{M}_O(P, Q, \theta) \sim \left(P(1)Q'(1) + \frac{1}{\theta}P'(1)Q(1) \right)^2 + \frac{1}{\theta} \int_0^1 \int_0^1 \left(\frac{1}{\theta}P''(x)Q(y) - \theta P(x)Q''(y) \right)^2 dx dy.$$

In this case we have $\mathcal{M}_O(x, 1, \theta) \sim \theta^{-2}$. It seems unexpected that $\mathcal{M}_O(x, 1, \theta) \rightarrow 1$ as $\theta \rightarrow \infty$. This may be related to the distribution of low-lying zeros of these L-functions (see [ILS]).

For the symplectic family, the mollifier has length \sqrt{X}^θ . The mollified mean value is

$$\mathcal{M}_{Sp}(P, Q, \theta) := \frac{1}{X^*} \sum_{c(f) \leq X} \left(Q \left(\frac{2}{\log X} \frac{d}{da} \right) \Phi_f \left(\frac{1}{2} + a \right) M \left(\frac{1}{2}, P \right) \right)^2 \Big|_{a=0}.$$

Applications of related mean values can be found in [S]. If Q is odd, then $\mathcal{M}_{Sp}(P, Q, \theta) = 0$. If Q is even, then for $\theta < 1$ we have

$$\mathcal{M}_{Sp}(P, Q, \theta) \sim \left(P(1)Q(1) + \frac{1}{\theta}P'(1)\widehat{Q}(1) \right)^2 + \frac{1}{\theta} \int_0^1 \int_0^1 \left(\frac{1}{\theta}P''(x)\widehat{Q}(y) - \theta P(x)Q'(y) \right)^2 dx dy,$$

where $\widehat{Q}(y) = \int_0^y Q(u) du$. We have $\mathcal{M}_{Sp}(x, 1, \theta) \sim (1 + \theta^{-1})^2$.

We end this discussion by pointing out the beautiful relationship

$$\mathcal{M}_{Sp}(P, Q', \theta) \sim \mathcal{M}_O(P, Q, \theta),$$

which is transparent in the above formulas. It was noted in [S] that this relationship is the source of the amazing “coincidence” of main results in [S] and [KM]. The same “coincidence” appears in the main result of [CGG], and it is plausible that there may be a similar explanation for that connection also.

4 Meromorphicity of g_λ

We prove the following theorem.

Theorem 4.1. The function $g_\lambda/\Gamma(1+B(\lambda))$ is meromorphic of order 2 in the whole complex plane. It never vanishes. Furthermore, for $k = 1, 2, \dots$,

(U) $g_{\lambda,u}/\Gamma(1+B_u(\lambda))$ has a pole of order $2k - 1$ at $\lambda = 1/2 - k$ and no other poles,

- (O) $g_{\lambda, O} / \Gamma(1 + B_O(\lambda))$ has a pole of order k at $\lambda = 1/2 - k$ and no other poles,
- (Sp) $g_{\lambda, Sp} / \Gamma(1 + B_{Sp}(\lambda))$ has a pole of order $k - 1$ at $\lambda = 1/2 - k$ and no other poles.

□

The zero and pole locations of $g_\lambda / \Gamma(1 + B(\lambda))$ indicate a connection with the Γ - and Γ_2 -function. By combining Theorem 4.1 with the asymptotic formulas for g_k given in Theorem 7.1, we obtain the following corollary.

Corollary 4.2. We have the following representations of g_λ in terms of the Γ -function and the Barnes-Vignéras double Γ -function:

$$\begin{aligned} \text{(U)} \quad & \frac{g_{\lambda, U}}{\Gamma(1 + B_U(\lambda))} = 2^{1/12} e^{3\zeta'(-1)} e^{-2\lambda\zeta'(0)} 2^{-2\lambda^2} \frac{\Gamma_2(\lambda + 1/2)^2}{\Gamma(\lambda + 1/2)}, \\ \text{(O)} \quad & \frac{g_{\lambda, O}}{\Gamma(1 + B_O(\lambda))} = 2^{-17/24} e^{(3/2)\zeta'(-1) + (1/2)\zeta'(0)} e^{-\lambda\zeta'(0)} 2^\lambda 2^{-(1/2)\lambda^2} \Gamma_2\left(\lambda + \frac{1}{2}\right), \\ \text{(Sp)} \quad & \frac{g_{\lambda, Sp}}{\Gamma(1 + B_{Sp}(\lambda))} = 2^{-5/24} e^{(3/2)\zeta'(-1) - (1/2)\zeta'(0)} e^{-\lambda\zeta'(0)} 2^{-\lambda} 2^{-(1/2)\lambda^2} \frac{\Gamma_2(\lambda + 1/2)}{\Gamma(\lambda + 1/2)}. \end{aligned}$$

In particular, $g_{\lambda+1, O} = 2^\lambda g_{\lambda, Sp}$, and

$$\frac{g_{\lambda, O}}{\Gamma(1 + B_O(\lambda))} \frac{g_{\lambda, Sp}}{\Gamma(1 + B_{Sp}(\lambda))} = 2^{\lambda^2 - 1} \frac{g_{\lambda, U}}{\Gamma(1 + B_U(\lambda))}. \quad \square$$

See [V] and [UN] for details about the double Γ -function.

Proof of Theorem 4.1. We illustrate with the case of $g_{\lambda, U}$; the other cases are similar. Choose J such that $1 + |2\lambda| < J < |4\lambda|$. Then for $j \geq J$ the real part of $j + 2\lambda$ is positive. We show that

$$\lim_{N \rightarrow \infty} \left(-\lambda^2 \log N + \sum_{j=J}^N \log \Gamma(j) - 2 \log \Gamma(j + \lambda) + \log \Gamma(j + 2\lambda) \right) \tag{4.1}$$

exists and is bounded by $|\lambda|^{2+\epsilon}$ for any $\epsilon > 0$. This assertion, together with well-known properties of the Γ -function, implies that $g_{\lambda, U} / \Gamma(1 + B_U(\lambda))$ is meromorphic of order at most 2, with zeros and poles as described. It follows that the order of $g_{\lambda, U} / \Gamma(1 + B_U(\lambda))$ is exactly 2 because the series $\sum \rho^{-r}$, where ρ runs through the set of poles, with multiplicity, has exponent of convergence $r = 2$.

Now we prove the above assertion. We need to estimate

$$\sum_{j=J}^N f_j(0) - 2f_j(\lambda) + f_j(2\lambda),$$

where $f_j(x) = \log \Gamma(x + j)$. By a mean value theorem,

$$f_j(0) - 2f_j(\lambda) + f_j(2\lambda) = \lambda^2 f_j''(\xi_j)$$

for some ξ_j between zero and 2λ .

To evaluate $f_j''(\xi_j)$, use the following well-known formula (see [R]), valid for $s \neq 0$ and not on the negative real axis,

$$\log \Gamma(s) = \left(s - \frac{1}{2}\right) \log s - s + \frac{1}{2} \log 2\pi + \frac{1}{2} \int_0^\infty \frac{\psi_2(u) + \frac{1}{6}}{(u + s)^2} du$$

with

$$\psi_2(u) = \{u\}^2 - \{u\}$$

and $\{u\} = u - [u]$. We have

$$\begin{aligned} f_j''(\xi) &= \frac{1}{\xi + j} - \frac{1}{2(\xi + j)^2} - 6 \int_0^\infty \frac{\psi_2(u)}{(u + j + \xi)^4} du \\ &= \frac{1}{j} + O\left(\frac{\xi}{j^2}\right). \end{aligned}$$

It follows that the limit in (4.1) exists and is bounded by $\lambda^2 \log J + O(\lambda/J) \ll 1 + \lambda^{2+\varepsilon}$. All the assertions are proved. ■

5 Integrality results

In this section k is a positive integer, and we are interested in integrality properties of g_k . By virtue of Corollary 4.2, these can be used to deduce rationality results for the double Γ -function. Denote by $v_p(n)$ the power of p dividing n .

Theorem 5.1. For all prime p we have $v_p(g_k) \geq 0$, so g_k is an integer. If $p > B(k)$, then $v_p(g_k) = 0$. If $p < B(k)$, then

$$v_p(g_k) = 0 \text{ iff } p^2 > B(k) \quad \text{and} \quad \begin{cases} k < p < k + \sqrt{p}, & (\text{U}) \\ k - \sqrt{k+p} < p < k + \sqrt{k+p} & (\text{O}) \end{cases}$$

with the Sp case following from the O case and the relationship $g_{k+1,O} = 2^k g_{k,Sp}$. □

Proof. We have the following useful formulas for g_k .

Lemma 5.2. For k a positive integer we have

$$g_{k,U} = B_U(k)! \frac{\left(\prod_{j=1}^{k-1} j!\right)^2}{2^{2k-1} \prod_{j=1}^{k-1} j!} = B_U(k)! 2^{k-k^2} \prod_{j=1}^{k-1} \frac{1}{(2j-1)!!(2j+1)!!},$$

$$g_{k,O} = B_O(k)! 2^{B_O(k)+k-1} \prod_{j=1}^{k-1} \frac{j!}{2j!} = B_O(k)! 2^{k-1} \prod_{j=1}^{k-1} \frac{1}{(2j-1)!!},$$

$$g_{k,Sp} = B_{Sp}(k)! 2^{B_{Sp}(k)} \prod_{j=1}^k \frac{j!}{2j!} = B_{Sp}(k)! \prod_{j=1}^k \frac{1}{(2j-1)!!}. \quad \square$$

Note that relationships between $g_{k,U}$, $g_{k,O}$, and $g_{k,Sp}$, as given at the end of Corollary 4.2, can also be obtained from the above formulas.

Proof. The proofs are similar, so we illustrate with the case of $g_{k,U}$. We begin with formula (2.1), first using the fact that k is an integer:

$$\begin{aligned} \frac{g_{k,U}}{B_U(k)!} &= N^{-k^2} \lim_{N \rightarrow \infty} \prod_{j=1}^N \prod_{m=1}^k \frac{j+2k-m}{j+k-m} \\ &= \lim_{N \rightarrow \infty} \prod_{m=1}^k N^{-k} \frac{\prod_{j=N-k+1}^N (j+2k-m)}{\prod_{j=1}^k (j+k-m)} \\ &= \lim_{N \rightarrow \infty} \prod_{m=1}^k \prod_{j=1}^k N^{-1} \frac{N-j+2k-m}{j+k-m} \\ &= \prod_{m=1}^k \prod_{j=1}^k \frac{1}{j+k-m} \\ &= \prod_{m=0}^{k-1} \prod_{j=1}^k \frac{1}{j+m} \\ &= \prod_{j=0}^{k-1} \frac{j!}{(j+k)!}. \end{aligned}$$

The last expression leads easily to the lemma. ■

From Lemma 5.2 it follows that $p \nmid g_k$ for the cases listed in Theorem 5.1. To prove the remaining statements in Theorem 5.1, we find a precise expression for $v_p(g_k)$.

We consider only the cases of $g_{k,U}$ and $g_{k,O}$. The remaining case follows immediately from the relationship $g_{k+1,O} = 2^k g_{k,S_p}$. Also, we consider only primes $p > 2$ because it is easy to show that g_k is divisible by a large power of 2. The main tool we need is the following lemma.

Lemma 5.3. We have

$$v_p \left(\prod_{j=1}^J j! \right) = \sum_{\ell=1}^{\infty} (J+1) \left[\frac{J}{p^\ell} \right] - \frac{p^\ell}{2} \left(\left[\frac{J}{p^\ell} \right]^2 + \left[\frac{J}{p^\ell} \right] \right)$$

and

$$v_p \left(\prod_{j=1}^J (2j-1)!! \right) = \sum_{\ell=1}^{\infty} \left(J + \frac{1}{2} \right) \left[\frac{2J-1}{p^\ell} \right]_2 - \frac{p^\ell}{2} \left[\frac{2J-1}{p^\ell} \right]_2^2,$$

where $[x]_2 = [(1/2)([x] + 1)]$. □

Proof. Regrouping the products gives

$$\prod_{j=1}^J j! = \prod_{j=1}^J j^{J-j+1} \quad \text{and} \quad \prod_{j=1}^J (2j-1)!! = \prod_{j=1}^J (2j-1)^{J-j+1}.$$

Thus,

$$v_p \left(\prod_{j=1}^J j! \right) = \sum_{\ell=1}^{\infty} \sum_{n=1}^{\lfloor J/p^\ell \rfloor} (J - np^\ell + 1)$$

and

$$v_p \left(\prod_{j=1}^J (2j-1)!! \right) = \sum_{\ell=1}^{\infty} \sum_{n=1}^{\lfloor (2J-1)/p^\ell \rfloor_2} \left(J + \frac{p^\ell + 1}{2} - np^\ell \right).$$

Evaluating the inner sums gives the lemma. ■

Combining Lemmas 5.2 and 5.3, we obtain the following expressions for $v_p(g_k)$.

Proposition 5.4. We have

$$v_p(g_{k,U}) = \sum_{\ell=1}^{\infty} \left(\left[\frac{k^2}{p^\ell} \right] + (2k - p^\ell) \left[\frac{k-1}{p^\ell} \right] + \left(\frac{p^\ell}{2} - 2k \right) \left[\frac{2k-1}{p^\ell} \right] - p^\ell \left[\frac{k-1}{p^\ell} \right]^2 + \frac{p^\ell}{2} \left[\frac{2k-1}{p^\ell} \right]^2 \right)$$

and

$$v_p(g_{k,O}) = \sum_{\ell=1}^{\infty} \left[\frac{\left(\frac{1}{2}\right)_{k(k-1)}}{p^\ell} \right] - \left(k - \frac{1}{2}\right) \left[\frac{2k-3}{p^\ell} \right]_2 + \frac{p^\ell}{2} \left[\frac{2k-3}{p^\ell} \right]_2^2. \quad \square$$

To complete the proof we must obtain lower bounds for the above expressions. The $g_{k,O}$ case is slightly simpler, so we handle it first.

Denote the summand in the second statement of Proposition 5.4 by $v_{p,\ell}(g_{k,O})$. Use θ to denote a number $0 \leq \theta < 1$. For example, $[x] = x - \theta$. We have

$$\begin{aligned} v_{p,\ell}(g_{k,O}) &= \left[\frac{\frac{1}{2}k(k-1)}{p^\ell} \right] - \left(k - \frac{1}{2}\right) \left[\frac{2k-3}{p^\ell} \right]_2 + \frac{p^\ell}{2} \left[\frac{2k-3}{p^\ell} \right]_2^2 \\ &= \frac{1}{2p^\ell} (k^2 - k) - \theta - \left(k - \frac{1}{2}\right) \left[\frac{2k-3}{p^\ell} \right]_2 + \frac{p^\ell}{2} \left[\frac{2k-3}{p^\ell} \right]_2^2 \\ &= \frac{1}{2p^\ell} \left(\left(k - p^\ell \left[\frac{2k-3}{p^\ell} \right]_2\right)^2 - \left(k - p^\ell \left[\frac{2k-3}{p^\ell} \right]_2\right) \right) - \theta \\ &> -1 \end{aligned} \tag{5.1}$$

because $M^2 - M \geq 0$ for integral M . Thus, $v_{p,\ell}(g_{k,O}) \geq 0$ for all p and ℓ , so $g_{k,O}$ is an integer. The remaining statements about $v_p(g_{k,O})$ can be obtained by considering the possibilities for $v_{p,\ell}(g_{k,O})$.

For the case of $g_{k,U}$, denote the summand in the first statement of Proposition 5.4 by $v_{p,\ell}(g_{k,U})$. There are two cases.

Case 1: $[(2k-1)/p^\ell] = 2[(k-1)/p^\ell]$. We have

$$\begin{aligned} v_{p,\ell}(g_{k,U}) &= \left[\frac{k^2}{p^\ell} \right] - 2k \left[\frac{k-1}{p^\ell} \right] + p^\ell \left[\frac{k-1}{p^\ell} \right]^2 \\ &= \frac{k^2}{p^\ell} - \theta - 2k \left[\frac{k-1}{p^\ell} \right] + p^\ell \left[\frac{k-1}{p^\ell} \right]^2 \\ &= p^{-\ell} \left(k - p^\ell \left[\frac{k-1}{p^\ell} \right] \right)^2 - \theta \\ &> -1. \end{aligned} \tag{5.2}$$

Thus, $v_{p,\ell}(g_{k,U}) \geq 0$.

Case 2: $[(2k - 1)/p^\ell] = 2[(k - 1)/p^\ell] + 1$. We have

$$\begin{aligned}
 v_{p,\ell}(g_{k,u}) &= \left[\frac{k^2}{p^\ell} \right] + (2p^\ell - 2k) \left[\frac{k-1}{p^\ell} \right] + p^\ell \left[\frac{k-1}{p^\ell} \right]^2 + p^\ell - 2k \\
 &= \frac{k^2}{p^\ell} - \theta + (2p^\ell - 2k) \left[\frac{k-1}{p^\ell} \right] + p^\ell \left[\frac{k-1}{p^\ell} \right]^2 + p^\ell - 2k \\
 &= p^{-\ell} \left(k - p^\ell - p^\ell \left[\frac{k-1}{p^\ell} \right] \right)^2 - \theta. \\
 &> -1.
 \end{aligned} \tag{5.3}$$

Again, $v_{p,\ell}(g_{k,u}) \geq 0$.

This covers all cases because for integral m, n , we have

$$\frac{n}{m} - 1 + \frac{1}{m} \leq \left[\frac{n}{m} \right] \leq \frac{n}{m} + 1 - \frac{1}{m}.$$

This proves that $g_{k,u}$ is an integer. The remaining statements can be proved by letting $\ell = 1$ and by considering the possible cases. This completes the proof of Theorem 5.1. ■

6 The function c_p

In this section we study the prime factorization of g_k as $k \rightarrow \infty$. We find that $v_p(g_k)$ is described by an interesting self-similar function.

Theorem 6.1. Define

$$c_p(x) = x^{-1} \sum_{\ell=-\infty}^{\infty} p^{-\ell} \|p^\ell x\|^2,$$

where $\|x\|$ is the distance from x to the nearest integer. Let $x > 0$, and put $k_j = [p^j x]$. Then as $j \rightarrow \infty$,

$$v_p(g_{k_j,u}) \sim k_j c_p(x) \quad \text{and} \quad v_p(g_{k_j,o}) \sim v_p(g_{k_j,sp}) \sim \frac{1}{2} k_j c_p(x). \quad \square$$

We leave the exercise of showing that $c_2(x) = 1$ to the reader. For the remainder of this section, we assume $p \geq 3$. We have some graphs in Figure 2.

Note that as $p \rightarrow \infty$, $c_p(x)$ approaches $x^{-1} \|x\|$, uniformly for $\delta \leq x \leq \delta^{-1}$.

From the graphs it appears that each $c_p(x)$ is not differentiable. Much more is actually true. At each rational point a/b , either the function $c_p(x)$ is self-similar, or it has a cusp, or it has a vertical tangent. This is made precise in Theorem 6.2.




Figure 2

Let $[[n]]$ denote the absolute least residue of $n \pmod b$; that is, $n \equiv [[n]] \pmod b$ and $-b/2 < [[n]] \leq b/2$.

Theorem 6.2. Suppose p , a , and b are pairwise coprime, and suppose $p^r \equiv 1 \pmod b$, with $r > 0$. Let $f(x) = ||x||^2$.

If $\sum_{j=0}^{r-1} [[ap^j]] = 0$, then $c_p(x)$ has the following self-similarity property at $x = a/b$:

$$\lim_{\substack{m \rightarrow \infty \\ m \equiv m_0 \pmod r}} \frac{c_p\left(\frac{a}{b} + p^{-m}x\right) - c_p\left(\frac{a}{b}\right)}{p^{-m}x} = \frac{x}{p-1} + \sum_{\ell=1-m_0}^{\infty} f'\left(p^{-\ell} \frac{a}{b}\right) + x^{-1} \sum_{\ell=0}^{\infty} p^{-\ell} \left(f\left(p^{\ell+m_0} \frac{a}{b} + p^{\ell}x\right) - f\left(p^{\ell+m_0} \frac{a}{b}\right) \right).$$

Note that this holds if $b = 1$ or more generally if $b \neq 2$ and $p^n \equiv -1 \pmod b$ for some n .

Suppose the above condition does not hold. In the case $b = 2$ we have

$$c_p\left(\frac{a}{2} + p^{-m}x\right) c_p = \left(\frac{a}{2}\right) - mp^{-m}|x| + O(p^{-m}x),$$

so c_p has a cusp at $a/2$. For all other b there is a nonzero constant k so that

$$c_p\left(\frac{a}{b} + p^{-m}x\right) c_p = \left(\frac{a}{b}\right) + kmp^{-m}x + O(p^{-m}x),$$

so c_p has a vertical tangent at a/b . □

Note that if $(p, b) \neq 1$, then one can use the relation $c_p(x) = c_p(p^jx)$ and then apply the theorem at $p^j a/b$.

We present some graphs of $c_p(x)$ near rational points a/b , where they are self-similar. The simplest case is when $b = 1$, for then $c_p(x)$ is self-similar under scaling

by $1/p$. In Figure 3, we give the examples of $c_3(x)$ near $x = 8$ and $c_7(x)$ near $x = 3$. In each graph the x -axis is interpreted as extending $1/p^m$ on either side of the central point, for any large m , and the vertical scale is also on the order of $1/p^m$.




Figure 3

Our last example is $c_5(x)$ near $x = 3/13$. Since 5 has order 4 mod 13 and since $5^2 \equiv -1 \pmod{13}$, the function is self-similar on rescaling by $1/p^4$. Thus, we have four possible pictures. In the graphs of Figure 4, the x -axis is interpreted as extending $1/p^m$ on either side of the central point, with $m \equiv i \pmod{4}$ for the graph in the i th quadrant. These graphs should be read counterclockwise, with each successive graph being the middle $1/5$ th of the previous graph.

Note that $c_5(3/13) = 23/72$. It is not difficult to evaluate $c_p(a/b)$ for any particular a/b , and in addition one sees that $c_p(a/b)$ is rational.

The self-similarity properties of $c_p(x)$ follow from the formula for $c_p(x)$ in Theorem 6.1. We have the following proposition.

Proposition 6.3. Let f be continuous on \mathbb{R} , periodic with period 1, and twice continuously differentiable except at finitely many points (mod 1) at which it is twice differentiable from both the left and the right. Also, suppose $f(x) \ll x^2$ and $f'(x) \ll |x|$ as $x \rightarrow 0$. Define

$$d(x) = \sum_{\ell=-\infty}^{\infty} p^{-\ell} f(p^\ell x).$$

Then $d(x)$ is continuous on \mathbb{R} and satisfies $d(px) = pd(x)$. Also, suppose $p^r \equiv 1 \pmod{b}$ and $m \equiv m_0 \pmod{r}$, with $m_0 \geq 0$. If x is sufficiently small, then

$$d\left(\frac{a}{b} + p^{-m}x\right) - d\left(\frac{a}{b}\right) = p^{-m}x \sum_{\ell=1-m_0}^{\infty} f'_\pm\left(p^{-\ell}\frac{a}{b}\right)$$

$$\begin{aligned}
 &+ p^{-m} \sum_{\ell=0}^{\infty} p^{-\ell} \left(f \left(p^{\ell+m_0} \frac{a}{b} + p^{\ell} x \right) - f \left(p^{\ell+m_0} \frac{a}{b} \right) \right) \\
 &+ (m - m_0) p^{-m} x \frac{1}{r} \sum_{j=0}^{r-1} f'_{\pm} \left(p^j \frac{a}{b} \right) \\
 &+ \frac{1}{2} p^{-2m} x^2 \sum_{\ell=0}^{m-1} p^{\ell} f''_{\pm} \left(p^{\ell} \frac{a}{b} + p^{\ell-m} \xi \right) + O(p^{-2m}),
 \end{aligned}$$

where $\xi = \xi_{\ell}$ is between zero and x . Here the \pm refers to a left- or right-derivative and has the same sign as x . □

Note that the condition that x be sufficiently small can be made explicit if one specifies the points at which f is not smooth.

$c_5(x)$ near $x = 3/13$




Figure 4

Proof of Theorem 6.2. Apply Proposition 6.3 with $f(x) = \|x\|^2$. We have

$$f'_{\pm} \left(\frac{a}{b} \right) = \begin{cases} f' \left(\frac{a}{b} \right) & \text{if } b \neq 2, \\ \mp 1 & \text{if } b = 2, \end{cases}$$

and $f''_{\pm}(a/b) = 2$ for all a/b . This makes the sum over f''_{\pm} explicit. For the second sum

over f'_\pm note that

$$\sum_{j=0}^{r-1} [[ap^j]] = \frac{b}{2} \sum_{j=0}^{r-1} f' \left(p^j \frac{a}{b} \right).$$

This proves Theorem 6.2. ■

It remains to prove Theorem 6.1 and Proposition 6.3.

Proof of Proposition 6.3. The continuity of $d(x)$ follows from the Weierstrass M-test, and the functional equation $d(px) = pd(x)$ follows by changing the summation index.

For the remaining properties, write

$$d\left(\frac{a}{b}\right) - d\left(\frac{a}{b} + p^{-m}x\right) = A + B + C,$$

where we used the definition of d and split the resulting sum into three pieces: $0 \leq \ell < m$, $m \leq \ell < \infty$, and $\ell < 0$.

We have

$$\begin{aligned} B &= \sum_{\ell=m}^{\infty} p^{-\ell} \left(f\left(p^\ell \frac{a}{b} + p^{\ell-m}x\right) - f\left(p^\ell \frac{a}{b}\right) \right) \\ &= p^{-m} \sum_{\ell=0}^{\infty} p^{-\ell} \left(f\left(p^{\ell+m} \frac{a}{b} + p^\ell x\right) - f\left(p^{\ell+m} \frac{a}{b}\right) \right) \\ &= p^{-m} \sum_{\ell=0}^{\infty} p^{-\ell} \left(f\left(p^{\ell+m_0} \frac{a}{b} + p^\ell x\right) - f\left(p^{\ell+m_0} \frac{a}{b}\right) \right). \end{aligned}$$

We did a change of variable; then we used the assumption $p^r \equiv 1 \pmod{b}$, the periodicity of f , and the fact that $m_0, \ell \geq 0$.

Next,

$$\begin{aligned} C &= \sum_{\ell=1}^{\infty} p^\ell \left(f\left(p^{-\ell} \frac{a}{b} + p^{-\ell-m}x\right) - f\left(p^{-\ell} \frac{a}{b}\right) \right) \\ &= \sum_{\ell=1}^{\infty} p^\ell \left(f'_\pm \left(p^{-\ell} \frac{a}{b} \right) p^{-\ell-m}x + O\left(p^{-2\ell-2m} f''_\pm \left(p^{-\ell} \frac{a}{b} \right) \right) \right) \\ &= p^{-m}x \sum_{\ell=1}^{\infty} f'_\pm \left(p^{-\ell} \frac{a}{b} \right) + O(p^{-2m}). \end{aligned}$$

We used Taylor's theorem and the fact that f'' is bounded.

Next,

$$\begin{aligned} A &= \sum_{\ell=0}^{m-1} p^{-\ell} \left(f\left(p^\ell \frac{a}{b} + p^{\ell-m} x\right) - f\left(p^\ell \frac{a}{b}\right) \right) \\ &= \sum_{\ell=0}^{m-1} p^{-\ell} \left(f'_\pm\left(p^\ell \frac{a}{b}\right) p^{\ell-m} x + \frac{1}{2} f''_\pm\left(p^\ell \frac{a}{b} + p^{\ell-m} \xi\right) p^{2\ell-2m} x^2 \right) \\ &= p^{-m} x \sum_{\ell=0}^{m-1} f'_\pm\left(p^\ell \frac{a}{b}\right) + \frac{1}{2} x^2 p^{-2m} \sum_{\ell=0}^{m-1} p^\ell f''_\pm\left(p^\ell \frac{a}{b} + p^{\ell-m} \xi\right), \end{aligned}$$

where $\xi = \xi_\ell$ is between zero and x . This is valid if x is sufficiently small, in terms of a/b , p , and f . Now,

$$\begin{aligned} \sum_{\ell=0}^{m-1} f'_\pm\left(p^\ell \frac{a}{b}\right) &= \sum_{j=0}^{r-1} \sum_{\ell=0}^{(m-m_0)/r-1} f'_\pm\left(p^{r\ell+j} \frac{a}{b}\right) + \sum_{j=0}^{m_0-1} f'_\pm\left(p^{m-m_0+j} \frac{a}{b}\right) \\ &= \sum_{j=0}^{r-1} \sum_{\ell=0}^{(m-m_0)/r-1} f'_\pm\left(p^j \frac{a}{b}\right) + \sum_{j=0}^{m_0-1} f'_\pm\left(p^j \frac{a}{b}\right) \\ &= \frac{m-m_0}{r} \sum_{j=0}^{r-1} f'_\pm\left(p^j \frac{a}{b}\right) + \sum_{j=0}^{m_0-1} f'_\pm\left(p^j \frac{a}{b}\right). \end{aligned}$$

This completes the proof of Proposition 6.3. ■

Proof of Theorem 6.1. First we consider the unitary case. We may unify formulas (5.2) and (5.3) to have

$$v_p(g_{k,U}) = \sum_{\ell=1}^{\infty} p^{-\ell} \left(k - p^\ell \left(\left[\frac{2k-1}{p^\ell} \right] - \left[\frac{k-1}{p^\ell} \right] \right) \right)^2 + O(\log k).$$

We have

$$\begin{aligned} \lim_{j \rightarrow \infty} \frac{v_p(g_{[p^j x], U})}{p^j} &= \lim_{j \rightarrow \infty} \frac{1}{p^j} \sum_{\ell=1}^{\infty} p^{-\ell} \left([p^j x] - p^\ell \left(\left[\frac{2[p^j x]-1}{p^\ell} \right] - \left[\frac{[p^j x]-1}{p^\ell} \right] \right) \right)^2 \\ &= \lim_{j \rightarrow \infty} \frac{1}{p^j} \sum_{\ell=1}^{\infty} p^{-\ell} \left(p^j x - \theta_j - p^\ell \left(\left[\frac{2p^j x - 2\theta_j - 1}{p^\ell} \right] - \left[\frac{p^j x - 1}{p^\ell} \right] \right) \right)^2 \\ &= \lim_{j \rightarrow \infty} \frac{1}{p^j} \sum_{\ell=1}^{\infty} p^\ell \left(p^{j-\ell} x - [2p^{j-\ell} x - (2\theta_j + 1)p^{-\ell}] + [p^{j-\ell} x - p^{-\ell}] \right)^2 \\ &= \lim_{j \rightarrow \infty} \sum_{\ell=-\infty}^{j-1} p^{-\ell} \left(p^\ell x - [2p^\ell x - (2\theta_j + 1)p^{\ell-j}] + [p^\ell x - p^{\ell-j}] \right)^2. \end{aligned}$$

We first used $[p^j x] = p^j x - \theta_j$, with $0 \leq \theta_j < 1$, and $[[x]/n] = [x/n]$. We eliminated the first θ_j by multiplying out and verifying that the other terms make no contribution. Then we changed variables $\ell \mapsto j - \ell$.

Let

$$a_{p,j}(x) = \lim_{\ell \rightarrow \infty} \sum_{\ell=-\infty}^{j-1} p^{-\ell} \left(p^\ell x - [2p^\ell x - (2\theta_j + 1)p^{\ell-j}] + [p^\ell x - p^{\ell-j}] \right)^2$$

denote the expression inside the above limit.

Let $[x]$ denote the largest integer strictly smaller than x . Note that

$$[t - \delta] = [t] \quad \text{if } 0 < \delta < t - [t].$$

Also, there is always such a δ because $t \neq [t]$ for all t . We use this to simplify $a_{p,j}(x)$.

Let $\varepsilon > 0$ be given, and choose $L > 0$ so that $p^{-L} < \varepsilon$. Then choose $\delta > 0$ such that

$$[2p^\ell x - \delta] = [2p^\ell x] \quad \text{and} \quad [p^\ell x - \delta] = [p^\ell x]$$

for $1 \leq \ell \leq L$. Now choose J so that $3p^{L-J} < \delta$. If $j \geq J$, we have

$$\begin{aligned} a_{p,j}(x) &= \sum_{\ell=-\infty}^L p^{-\ell} \left(p^\ell x - [2p^\ell x] + [p^\ell x] \right)^2 \\ &\quad + \sum_{\ell=L+1}^{j-1} p^{-\ell} \left(p^\ell x - [2p^\ell x - (2\theta_j + 1)p^{\ell-j}] + [p^\ell x - p^{\ell-j}] \right)^2 \\ &= \sum_{\ell=-\infty}^{\infty} p^{-\ell} \left(p^\ell x - [2p^\ell x] + [p^\ell x] \right)^2 + O\left(\sum_{\ell=L+1}^{\infty} p^{-\ell} + \sum_{\ell=L+1}^{J-1} p^{-\ell} \right) \\ &= \sum_{\ell=-\infty}^{\infty} p^{-\ell} \left(p^\ell x - [2p^\ell x] + [p^\ell x] \right)^2 + O(p^{-L}), \end{aligned}$$

and we were given that $p^{-L} < \varepsilon$. Thus,

$$\lim_{j \rightarrow \infty} a_{p,j}(x) = \sum_{\ell=-\infty}^{\infty} p^{-\ell} \left(p^\ell x - [2p^\ell x] + [p^\ell x] \right)^2.$$

To finish the proof, note that $(x - [2x] + [x])^2 = (x - [2x] + [x])^2 = \|x\|^2$.

For the orthogonal case, a similar calculation shows

$$\begin{aligned} \lim_{j \rightarrow \infty} \frac{v_p(g_{[p^j x], 0})}{p^j} &= \frac{1}{2} \sum_{\ell=-\infty}^{\infty} p^{-\ell} \left(p^\ell x - [2p^\ell x]_2 \right)^2 \\ &= \frac{1}{2} x c_p(x), \end{aligned}$$

since $(x - [2x]_2)^2 = \|x\|^2$. The same holds in the symplectic case because $g_{k+1, \mathcal{O}} = 2^k g_{k, \text{Sp}}$. This completes the proof of Theorem 6.1. \blacksquare

7 Asymptotics

In this section we determine the asymptotic behavior of g_k for large (integer) k . We have the following theorem.

Theorem 7.1. As $k \rightarrow \infty$,

$$\begin{aligned}
 \log g_{k, \mathcal{U}} &= \log B_{\mathcal{U}}(k)! - k^2 \log k + \left(\frac{3}{2} - 2 \log 2\right) k^2 - \frac{1}{12} \log k \\
 &\quad + \frac{1}{12} \log 2 + \zeta'(-1) + O(k^{-1}) \\
 &= k^2 \log k + \left(\frac{1}{2} - 2 \log 2\right) k^2 + \frac{11}{12} \log k \\
 &\quad + \frac{1}{12} \log 2 - \zeta'(0) + \zeta'(-1) + O(k^{-1}), \\
 \log g_{k, \mathcal{O}} &= \log B_{\mathcal{O}}(k)! - \frac{1}{2} k^2 \log k + \left(\frac{3}{4} - \frac{1}{2} \log 2\right) k^2 + \frac{1}{2} k \log k + \left(\log 2 - \frac{1}{2}\right) k \\
 &\quad - \frac{1}{24} \log k - \frac{17}{24} \log 2 + \frac{1}{2} \zeta'(-1) + O(k^{-1}) \\
 &= \frac{1}{2} k^2 \log k + \left(\frac{1}{4} - \log 2\right) k^2 - \frac{1}{2} k \log k + \left(\frac{3}{2} \log 2 - \frac{1}{2}\right) k + \frac{23}{24} \log k \\
 &\quad - \frac{29}{24} \log 2 + \frac{1}{4} - \zeta'(0) + \frac{1}{2} \zeta'(-1) + O(k^{-1}), \\
 \log g_{k, \text{Sp}} &= \log B_{\text{Sp}}(k)! - \frac{1}{2} k^2 \log k + \left(\frac{3}{4} - \frac{1}{2} \log 2\right) k^2 - \frac{1}{2} k \log k + \left(\frac{1}{2} - \log 2\right) k \\
 &\quad - \frac{1}{24} \log k - \frac{5}{24} \log 2 + \frac{1}{2} \zeta'(-1) + O(k^{-1}) \\
 &= \frac{1}{2} k^2 \log k + \left(\frac{1}{4} - \log 2\right) k^2 + \frac{1}{2} k \log k + \left(\frac{1}{2} - \frac{3}{2} \log 2\right) k + \frac{23}{24} \log k \\
 &\quad - \frac{17}{24} \log 2 + \frac{1}{4} - \zeta'(0) + \frac{1}{2} \zeta'(-1) + O(k^{-1}). \quad \square
 \end{aligned}$$

Combining these with the asymptotics of the Γ - and Γ_2 -function (see [V] and [UN]), we obtain the expressions for g_λ in terms of Γ_2 given in Corollary 4.2.

Proof. To prove Theorem 7.1, combine the expressions for g_λ given in Lemma 5.1 with the elementary formulas

$$\sum_{j=1}^n \log j! = (n+1) \sum_{j=1}^n \log j - \sum_{j=1}^n j \log j$$

and

$$\begin{aligned} \sum_{j=1}^n \log 2j! &= \frac{1}{2}n(n+1)\log 2 + (n+1)\sum_{j=1}^n \log j + (n+1)\sum_{j=1}^n \log(2j-1) \\ &\quad - \sum_{j=1}^n j \log j - \sum_{j=1}^n j \log(2j-1). \end{aligned}$$

The formulas in the following lemma are sufficient to complete the calculations.

Lemma 7.2. As $n \rightarrow \infty$ we have

$$\begin{aligned} \sum_{j=1}^n \log j &= n \log n - n + \frac{1}{2} \log n - \zeta'(0) + \frac{1}{12n} + O(n^{-2}), \\ \sum_{j=1}^n \log(2j-1) &= n \log 2n - n + \frac{1}{2} \log 2 - \frac{1}{24n} + O(n^{-2}), \\ \sum_{j=1}^n j \log j &= \frac{1}{2}n^2 \log \frac{1}{4}n^2 + \frac{1}{2}n \log n + \frac{1}{12} \log n + \frac{1}{12} - \zeta'(-1) + O(n^{-1}), \\ \sum_{j=1}^n j \log(2j-1) &= \frac{1}{2}n^2 \log 2n - \frac{1}{4}n^2 + \frac{1}{2}n \log 2n - \frac{1}{2}n - \frac{1}{24} \log n \\ &\quad + \frac{7}{24} \log 2 - \frac{1}{24} + \frac{1}{2} \zeta'(-1) + O(n^{-1}). \quad \square \end{aligned}$$

The first formula in Lemma 7.2 is Stirling's formula, and each expression can be derived from the Euler-Maclaurin summation formula (see [R]). This completes the proof of Theorem 7.1. ■

Acknowledgments

This research was supported by the American Institute of Mathematics. Conrey's work was supported in part by a grant from the National Science Foundation. We thank Keating, Rudnick, Sarnak, Selberg, and Snaith for helpful conversations.

References

- [BK] E. B. Bogomolny and J. P. Keating, *Random matrix theory and the Riemann zeros, II: n-point correlations*, *Nonlinearity* **9** (1996), 911–935.
- [BH] E. Brézin and S. Hikami, *Characteristic polynomials of random matrices*, preprint, 1999.
- [C] J. B. Conrey, *More than two fifths of the zeros of the Riemann zeta function are on the critical line*, *J. Reine Angew. Math.* **399** (1989), 1–26.

- [CG1] J. B. Conrey and A. Ghosh, *On mean values of the zeta-function*, *Mathematika* **31** (1984), 159–161.
- [CG2] ———, “Mean values of the Riemann zeta-function, III” in *Proceedings of the Amalfi Conference on Analytic Number Theory (Maiori, 1989)*, Univ. Salerno, Salerno, 1992, 35–59.
- [CG3] ———, *A conjecture for the sixth power moment of the Riemann zeta-function*, *Internat. Math. Res. Notices* **1998**, 775–780.
- [CGG] J. B. Conrey, A. Ghosh, and S. M. Gonek, *Simple zeros of the Riemann zeta-function*, *Proc. London Math. Soc.* (3) **76** (1998), 497–522.
- [CGo] J. B. Conrey and S. M. Gonek, *High moments of the Riemann zeta-function*, preprint, 1999.
- [D] W. Duke, *The critical order of vanishing of automorphic L-functions with large level*, *Invent. Math.* **119** (1995), 165–174.
- [DFI] W. Duke, J. Friedlander, and H. Iwaniec, *Bounds for automorphic L-functions, II*, *Invent. Math.* **115** (1994), 219–239.
- [G] S. M. Gonek, *On negative moments of the Riemann zeta-function*, *Mathematika* **36** (1989), 71–88.
- [HB] D. R. Heath-Brown, *Fractional moments of the Riemann zeta-function, II*, *Quart. J. Math. Oxford Ser. (2)* **44** (1993), 185–197.
- [HL] G. H. Hardy and J. E. Littlewood, *Contributions to the theory of the Riemann zeta-function and the theory of the distribution of primes*, *Acta Math.* **41** (1918), 119–196.
- [I] A. E. Ingham, *Mean-value theorems in the theory of the Riemann zeta-function*, *Proc. London Math. Soc.* (2) **27** (1926), 273–300
- [ILS] H. Iwaniec, W. Luo, and P. Sarnak, *Low lying zeros of families of L-functions*, preprint, 1998.
- [IS] H. Iwaniec and P. Sarnak, *Mean values of L-functions and the Landau-Siegel zero*, preprint, 1997.
- [J] M. Jutila, *On the mean value of $L(1/2, \chi)$ for real characters*, *Analysis* **1** (1981), 149–161
- [KS1] N. M. Katz and P. Sarnak, *Zeros of zeta functions and symmetry*, *Bull. Amer. Math. Soc. (N.S.)* **36** (1999), 1–26.
- [KS2] ———, *Random Matrices, Frobenius Eigenvalues, and Monodromy*, *Amer. Math. Soc. Colloq. Publ.* **45**, Amer. Math. Soc., Providence, 1999.
- [KeSn] J. Keating and N. Snaith, *Random matrix theory and some zeta-function moments*, lecture at Erwin Schrödinger Institute, 1998, and personal communication, 1999.
- [KM] E. Kowalski and P. Michel, *A lower bound for the rank of $J_0(q)$* , preprint, 1999.
- [KMV] E. Kowalski, P. Michel, and J. VanderKam, *Nonvanishing of high derivatives of automorphic L-functions at the center of the critical strip*, preprint, 1999.
- [Lev] N. Levinson, *More than one third of zeros of Riemann’s zeta-function are on $\sigma = 1/2$* , *Adv. Math.* **13** (1974), 383–436.
- [M] H. L. Montgomery, “The pair correlation of zeros of the zeta function” in *Analytic Number Theory (St. Louis, Mo., 1972)*, *Proc. Sympos. Pure Math.* **24**, Amer. Math. Soc., Providence, 1973, 181–193.
- [MV] H. L. Montgomery and R. C. Vaughan, *The large sieve*, *Mathematika* **20** (1973), 119–134.
- [O] A. M. Odlyzko, *The 10^{20} zero of the Riemann zeta-function and 70 million of its neighbors*, preprint.

- [R] H. Rademacher, *Topics in Analytic Number Theory*, Grundlehren Math. Wiss. **169**, Springer, New York, 1973.
- [Ru] M. O. Rubinstein, *Evidence for a spectral interpretation of the zeros of L-function*, Ph.D. dissertation, Princeton Univ., Princeton, 1998.
- [Sa] P. Sarnak, *Determinants of Laplacians*, Comm. Math. Phys. **110** (1987), 113–120.
- [S] K. Soundararajan, *Nonvanishing of quadratic Dirichlet L-functions at $s = 1/2$* , preprint, 1999.
- [UN] K. Ueno and M. Nishizawa, *Multiple gamma functions and multiple q-gamma functions*, Publ. Res. Inst. Math. Sci. **33** (1997), 813–838
- [V] M.-F. Vignéras, "L'equation fonctionnelle de la fonction zêta de Selberg du groupe modulaire $PSL(2, \mathbb{Z})$ " in *Journées Arithmétiques de Luminy (Luminy, 1978)*, Astérisque **61**, Soc. Math. France, Paris, 1979, 235–249.

Conrey: American Institute of Mathematics, 360 Portage Avenue, Palo Alto, California 94306, USA;
conrey@aimath.org

Farmer: Department of Mathematics, Bucknell University, Lewisburg, Pennsylvania 17837, USA;
farmer@bucknell.edu